

# The Holocene

<http://hol.sagepub.com/>

---

## **Abrupt cooling repeatedly punctuated early-Holocene climate in eastern North America**

Juzhi Hou, Yongsong Huang, Bryan N Shuman, W Wyatt Oswald and David R Foster

*The Holocene* published online 30 November 2011

DOI: 10.1177/0959683611427329

The online version of this article can be found at:

<http://hol.sagepub.com/content/early/2011/11/28/0959683611427329>

---

Published by:



<http://www.sagepublications.com>

**Additional services and information for *The Holocene* can be found at:**

**Email Alerts:** <http://hol.sagepub.com/cgi/alerts>


**Subscriptions:** <http://hol.sagepub.com/subscriptions>

**Reprints:** <http://www.sagepub.com/journalsReprints.nav>


**Permissions:** <http://www.sagepub.com/journalsPermissions.nav>

>> **Proof** - Nov 30, 2011

[What is This?](#)



# Abrupt cooling repeatedly punctuated early-Holocene climate in eastern North America

The Holocene  
0(0) 1–5  
© The Author(s) 2011  
Reprints and permission:  
sagepub.co.uk/journalsPermissions.nav  
DOI: 10.1177/0959683611427329  
hol.sagepub.com  


Juzhi Hou,<sup>1,2</sup> Yongsong Huang,<sup>2</sup> Bryan N. Shuman,<sup>3</sup>  
W. Wyatt Oswald<sup>4</sup> and David R. Foster<sup>5</sup>

## Abstract

Climate proxy records and general circulation models suggest that Atlantic Meridional Overturning Circulation (AMOC) plays a key role for global climate changes. Paleocceanographic data document multiple episodes of prominent AMOC weakening during the early Holocene. However, proxy records at adjacent continents have not been demonstrated to fully capture the climate responses to multiple AMOC variation due to temporal resolution and/or the proxy sensitivity. Here we present decadal- to multidecadal-resolution hydrogen isotopic records of aquatic biomarkers from Blood Pond, Massachusetts during the early Holocene. Our data reveal a full series of prominent and abrupt cooling events centered on 10.6, 10.2, 9.5, 9.2, 8.8 and 8.4 ka. These abrupt climatic reversals coincide with key intervals of weakened AMOC, suggesting an apparent relationship between AMOC oscillations and the abrupt continental climate changes in northeastern North America. The noticeable connection implies that the AMOC variation did play an important role in the abrupt climate changes during the early Holocene. Our data also suggest that northeastern North America may experience significant climatic variations should the predicted major disturbance of AMOC occur in the coming century as a result of anthropogenic greenhouse gas emissions.

## Keywords

abrupt climate event, AMOC, compound-specific D/H ratios, early Holocene, northeastern North America

## Introduction

Abrupt climate changes exert significant impacts on natural ecosystems and human societies (Overpeck and Cole, 2006). One widely proposed mechanism for abrupt climate changes is perturbation of the Atlantic Meridional Overturning Circulation (AMOC) (Clark et al., 2002; Overpeck and Cole, 2006), which occurred numerous times from the Late Pleistocene to the early Holocene as a result of freshwater outbursts from proglacial lakes which originated from the melting Laurentide ice sheet (LIS) (Clark et al., 2001; Stouffer et al., 2006). The Younger Dryas (YD) and the 8.2 ka events are two of the well-known abrupt climate events attributed to slowdown in AMOC (Alley and Agustsdottir, 2005; Clark et al., 2001, 2002; Clarke et al., 2009; LeGrande et al., 2006). However, carbon isotopic ratios of benthic foraminifera and glacial landform mapping record multiple episodes of freshwater outbursts and AMOC changes from the Late Pleistocene to the early Holocene (Clark et al., 2001; Teller and Leverington, 2004). A few terrestrial and marine records suggest the influence of the AMOC on the decadal- to centennial-scale climate change in the North Atlantic and neighboring regions (Bamberg et al., 2010; Daley et al., 2009). Of particular significance is the absence of records of abrupt climate changes in a critical region – northeastern North America, which climate models indicate is highly sensitive to AMOC. The inconsistency between the proposed climate driver (AMOC) and climate records may result from (1) the fact that the existing climate records from northeastern North America cannot fully capture the relatively short scale (decadal to multidecadal) climate variability, either because of insufficient temporal resolution or inadequate proxy sensitivity; (2) regional climate sensitivity to AMOC differs under different climatic boundary conditions, with sensitivity being particularly high at the YD and 8.2 events. General

circulation model simulations predict substantial reductions in the strength of AMOC with the ongoing rapid rise in anthropogenic greenhouse gases (Schmidt et al., 2005), but inconsistency in the paleoclimate records of the North Atlantic and North America raises important questions about the climatic sensitivity to changes in AMOC. This inconsistency has fueled significant debates over the role of AMOC on abrupt climate variability, with some arguing a pivotal role (Rhines et al., 2008), whereas others a negligible one (Seager and Battisti, 2007).

Here, we present decadal-scale compound-specific hydrogen isotope records from Blood Pond, south-central Massachusetts, USA (Figures 1 and 2). The D/H ratios of behenic acid ( $\delta D_{BA}$ ), mainly produced by aquatic macrophytes at Blood Pond (Hou et al., 2006), record lake water isotope composition and hence precipitation isotopic ratios in a cool and wet region such as New England (Hou et al., 2007). We have previously demonstrated that the centennial- to millennial-scale  $\delta D_{BA}$  records from Blood Pond

<sup>1</sup>Institute of Tibetan Plateau Research, Chinese Academy of Sciences, China

<sup>2</sup>Brown University, USA

<sup>3</sup>University of Wyoming, USA

<sup>4</sup>Emerson College, USA


<sup>5</sup>Harvard University, USA

Received 8 December 2010; revised manuscript accepted 26 September 2011


## Corresponding author:

Juzhi Hou, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing 100083, China.

Email: houjz@itpcas.ac.cn



**Figure 1.** Map showing the location of Blood Pond and other sites discussed in the text. Three routes of freshwater outbursts are indicated: (A) Hudson Bay (8.2 ka event); (B) North Atlantic Ocean; (C) Gulf of Mexico. Key components of the Atlantic Meridional Overturning Circulation (AMOC) relevant to this study are also shown



**Figure 2.** Comparison of hydrogen isotope records from Blood Pond (71.961°W, 42.081°N) with other terrestrial records. (A) Inferred global terrestrial  $^{18}\text{O}/^{16}\text{O}$  fractionation ( $\Delta\epsilon_{\text{LAND}} \text{‰}$ ) based on  $\delta^{18}\text{O}$  of oxygen gas from the Siple Dome ice core, Antarctica (Xu and Liu, 2007). (B) Sediment grain size from Rocky Pond, Massachusetts (Kaser et al., 2004). (C) Temperature inferred from chironomid assemblages in Lake CF8, Baffin Island (Haeblerli and Gruber, 2009). (D) Stable hydrogen isotope records of  $\text{C}_{22}$   $n$ -alkanoic acid ( $\delta\text{D}_{\text{BA}}$ ) and inferred temperature variability. Chronology of the Blood Pond sediment core is defined by a total of 15 accelerator mass spectrometry radiocarbon dates (5 AMS dates for the study section in this study)

faithfully capture the regional climate history (Hou et al., 2006, 2007). The objectives of this study are: (1) to investigate the relationship between the decadal- to multidecadal-scale hydrogen isotope records during the early Holocene and the reconstructed freshwater outbursts from the LIS as well as the inferred AMOC changes; (2) to further evaluate the effect of AMOC variations on adjacent continental climate changes.

## Samples and methods


Blood Pond (42.081°N, 71.961°W, 212.1 m a.s.l.) is a kettle pond located in Massachusetts (Figure 1). The lake is mainly recharged by groundwater and precipitation, with a southerly overflow outlet. The sediment core was collected in 2001, and samples for isotope analyses were selected at 2 cm intervals. Chronology for the core is provided by AMS- $^{14}\text{C}$  dating of bulk organic matter which were converted to calendar years before present (cal. yr BP) using OxCal 3.9 (Hou et al., 2006). The sedimentation rate of the early Holocene (11–7 ka) at Blood Pond is  $\sim 1.3$  cm/10 yr, allowing for the evaluation of decadal- to centennial-scale climate variation.

The preparation of the samples and the measurement of hydrogen isotopes have been described previously (Hou et al., 2006). Briefly, lipids were extracted from freeze-dried sediment using an Accelerated Solvent Extractor 200 (Dionex). Acid fractions were isolated, methylated, and purified. A HP 6890 GC interfaced to a Finnigan Delta+ XL stable isotope spectrometer through a high-temperature pyrolysis reactor was used for hydrogen isotopic analysis. The precision ( $1\sigma$ ) of triplicate analyses was  $< \pm 2\%$ . The accuracy was routinely checked by measuring laboratory isotopic standards every six measurements. The  $\delta\text{D}$  value of the added methyl group was determined by acidifying and then methylating (along with the samples) the disodium salt of succinic acid with a predetermined  $\delta\text{D}$  value (using TC/EA-IRMS) (Huang et al., 2002).

## Results and discussion

### Early-Holocene climate revealed by Blood Pond $\delta\text{D}_{\text{BA}}$

Our new decadal to multidecadal  $\delta\text{D}_{\text{BA}}$  record during the early Holocene reveals six distinct negative isotopic excursions centered at 10.6, 10.2, 9.5, 9.2, 8.8 and 8.4 ka (Figure 2). The largest and longest of these isotopic excursions occurs around 9.2–9.3 ka with  $\sim 30\%$   $\delta\text{D}_{\text{BA}}$  shift, and two significant  $\delta\text{D}_{\text{BA}}$  shifts ( $\sim 15\%$ ) occur around 10.2 and 8.4 ka. The isotopic shift at 9.2 ka is, within the chronological uncertainties, concurrent with changes seen in Greenland ice core  $\delta^{18}\text{O}$  (Vinther et al., 2006), European lake sediment  $\delta^{18}\text{O}$  (von Grafenstein et al., 1999) and tree ring width (Spurk et al., 2002), Baffin Island Chironomid data (Kurek et al., 2004), East Asian and Arabian speleothem  $\delta^{18}\text{O}$  records (Dykoski et al., 2005; Fleitmann et al., 2008), and Alaskan lake biogenic silica (Hu et al., 2003). Recent data from isotopic ratios of oxygen gas preserved in the Siple dome ice core display a clear 9.2 ka shift (Severinghaus et al., 2009), strongly supporting a near-global 9.2 ka event (Fleitmann et al., 2008). The freshwater input responsible for the 9.2–9.3 ka cooling event might come from Lake Superior as lake level rapidly dropped for 45 m at the same time, which was delivered into the North Atlantic Ocean via Lake Huron–North Bay–Ottawa River–St Lawrence River valleys (Yu et al., 2010). Our results are also supported by regional paleoclimate data from New England (Figure 2). Cooling events in the early Holocene inferred by our isotopic records are closely associated with overall drier conditions as indicated by shifts in sediment facies (increase in grain sizes) at nearby Rocky and New Long Ponds, Massachusetts (Newby et al., 2009). A modern analog for such a cold-dry scenario is the AD 1962–1965 drought, which was the most severe historic drought in the northeastern USA (Namias, 1966). During AD 1962–1965 a band of cool sea-surface temperatures extending from Maine to Ireland severely reduced regional summer precipitation by drawing cold Arctic air masses into the region. These air masses also led to unusually low temperatures, especially during the winter. The cooler and drier climate around 9.2 ka, as demonstrated by  $\delta\text{D}_{\text{BA}}$  data, also coincides with a broad interval of elevated *Ambrosia* pollen reflecting



**Figure 3.** Comparison of Blood Pond  $\delta D_{BA}$  records with records inferring changes in the AMOC. (A) Freshwater release to North Atlantic Ocean (Teller and Leverington, 2004). (B) Detrended  $\Delta^{14}C$  from tree rings (Stuiver et al., 1998). (C)  $\delta^{13}C$  records from VM29-191 (Bond et al., 2001) and Blood Pond  $\delta D_{BA}$  records. The climate reversals at Blood Pond are strongly correlated with inferred AMOC changes, possibly induced by freshwater outbursts from Laurentide Ice Sheet

more open land in northeastern North America (Faison et al., 2006).

#### Correlation with AMOC variability

The abrupt climate reversals revealed at Blood Pond correspond strongly to the early-Holocene AMOC changes inferred from foraminifera  $\delta^{13}C$  values as well as detrended atmospheric  $\Delta^{14}C$  record from tree rings (Stuiver et al., 1998) (Figure 3). The benthic foraminifera  $\delta^{13}C$  values at site VM29-191 essentially reflect the convective overturning rate of AMOC, with lower  $\delta^{13}C$  implying reduced overturning (Bond et al., 1999; Clark et al., 2001). The definitive identification of a 10.2 ka climate excursion at Blood Pond, which coincides with the most severe lake low stand at Rocky Pond (Newby et al., 2009), is particularly important, because the 10.2 ka  $\delta^{13}C$  change at VM29-191 is as large as 0.6‰, accounting for ~70% amplitude of the  $\delta^{13}C$  excursions during the YD and suggesting substantial AMOC reductions. The increase in  $\Delta^{14}C$  at the abrupt climate reversals could also indicate AMOC slowdowns during the early Holocene (Clark et al., 2001). The fact that all the abrupt hydrogen isotopic excursions during the early Holocene at Blood Pond have counterparts within the  $\delta^{13}C$  and/or  $\Delta^{14}C$  data suggests a consistent, causal relationship between AMOC strength and climate variability in New England. Likewise, the evidence for cool, dry climate in New England when AMOC slowed is consistent with historic responses to North Atlantic conditions (Namias, 1966).


#### The 10.2 and 9.2 ka events

Blood Pond provides the first isotopic record from North America to register abrupt climatic changes at 9.2 and 10.2 ka. However, directly applying the  $\delta D_{BA}$ -temperature relationship ( $\delta D_{BA} = 4.3T - 208.4$ ) (Hou et al., 2006) established from a transect of eastern North American lakes yields an unexpectedly large

maximum temperature drop of nearly  $\sim 7^{\circ}C$  around 9.2 ka, only slightly smaller than that estimated for the YD (Hou et al., 2007). It is thus unlikely that the isotopic shift at 9.2 kyr BP represents mean annual temperature change alone. Precipitation seasonality must have also changed, with an increase in the ratio of cool to warm season precipitation. This is consistent with early-Holocene lake-level low stands in New England, which have been attributed to reduced summer precipitation and relatively high winter precipitation (Shuman and Donnelly, 2006). The modern-day precipitation  $\delta D$  values at Blood Pond are  $\sim -40$ ‰ for June to August and  $\sim -82$ ‰ for November to January, with precipitation amounts evenly distributed throughout the year (Bowen and Revenaugh, 2003). A 20% decrease in summer relative to winter precipitation would produce a  $\sim 13$ ‰ reduction in the  $\delta D$  values of mean annual precipitation. Likewise, the historic analog to these events produced a severe reduction in summer precipitation (Namias, 1966). If temperature declines were primarily a wintertime feature, as models have inferred for AMOC induced temperature decreases (Seager and Battisti, 2007), the lower  $\delta D$  values of the colder winter precipitation will further reduce annual mean precipitation D/H ratios. Therefore, temperature derived from  $\delta D_{BA}$  changes should be more appropriately defined as precipitation-weighted temperature (PWT), taking into consideration seasonal shifts in precipitation and temperature. Moreover, the large size of the Laurentide Ice Sheet at 9.2 ka (two times the modern Greenland) (Dyke and Prest, 1987) and relative proximity to our study site may have been particularly important for the very different seasonality in precipitation and unusually large 9.2 ka isotope excursion.

#### The 8.2 ka and other events

In addition to 9.2 and 10.2 ka climatic reversals, the  $\delta D_{BA}$  record also shows a clear abrupt decrease from 8.45 to 8.3 kyr BP (Figure 2), with a  $\sim 3$ – $4^{\circ}C$  decline in PWT change. This temperature shift slightly precedes the well-known 8.2 ka event in the ice cores and speleothems by about 200 years (Alley and Agustsdottir, 2005). Currently, it is unclear whether the timing discrepancy results from chronological inaccuracies or actual difference in timing of the continental climatic response. We adopted a linear interpolation between the neighboring  $^{14}C$  ages to assign the ages of individual samples during the early Holocene. The sedimentation rate might have changed between the two neighboring  $^{14}C$  ages, which would cause the small age offsets. However, the timing of the negative  $\delta D_{BA}$  shift between 8.45 and 8.3 kyr BP is consistent with climatic and ecological records from northeastern North America (Kurek et al., 2004; Lutz et al., 2007; Shuman et al., 2002). Ellison et al. (2006) suggest the presence of two distinct cooling events centered at 8.5 and 8.1 kyr BP, respectively, in the subpolar North Atlantic Ocean (Ellison et al., 2006). The chronological uncertainties of the Blood Pond sediment core make it difficult to assign the  $\delta D_{BA}$  shift between 8.45 and 8.3 kyr BP to one of these two events. Additionally, the precipitation seasonality during the 8.2 ka event may differ from that during the 9.2 ka event because of the final collapse of Laurentide Ice Sheet. The collapse of the LIS may have resulted in a more evenly distributed precipitation throughout the year, which may explain the relatively small  $\delta D_{BA}$  shift during the 8.2 ka event. At 10.6 ka,  $\delta D_{BA}$  decreased by  $\sim 10$ ‰ with a duration of about a century, which corresponds to a similar  $\delta^{13}C$  variation (Clark et al., 2001). In addition to the above four clearly identified century-scale climatic reversals, we also found two relatively short-lived  $\delta D_{BA}$  excursions at 8.8 and 9.5 ka, lasting only a couple of decades. Of these brief excursions, only the one at 8.8 ka has a corresponding carbon isotopic excursion in VM29-191. Definitive identification of abrupt climate events of such short duration will require higher resolution isotopic records.



**Figure 4.** Compilation of selected records showing the abrupt climate reversals during the early Holocene at 8.2 ka, 8.8 ka, 9.2 ka, 9.6 ka, 10.2 ka and 10.6 ka. (A) Grey scale from Cariaco Basin (Hughen et al., 2000). (B) Ti concentration from Huguangyan Maar Lake (Yancheva et al., 2007). (C) Residue on ignition (ROI) from NAUJHI (Willemse and Tornqvist, 1999). (D)  $\delta^{18}\text{O}$  records from Lake Ammersee (von Grafenstein et al., 1999). (E)  $\delta^{18}\text{O}$  records from NGRIP (Vinther et al., 2006). (F)  $\delta^{18}\text{O}$  records from Gunnug Buda cave. (G)  $\delta^{18}\text{O}$  records from Qunf Cave (Fleitmann et al., 2008). (H)  $\delta^{18}\text{O}$  records from Dongge Cave (Dykoski et al., 2005). (I) Blood Pond  $\delta\text{D}_{\text{BA}}$  records

### Comparison with global climate records


The abrupt climate reversals revealed in the Blood Pond records can also be tentatively found in other high-resolution proxy records of climate around the globe (Figures 4 and 5). Other than the Greenland ice cores, most other quantitative climate records are derived from speleothem  $\delta^{18}\text{O}$  values. To improve the comparability with our hydrogen isotopic records, we converted the  $\delta\text{D}_{\text{BA}}$  excursions at Blood Pond into oxygen isotopic variations using the slope of the global meteoric water line (i.e.  $\delta\text{D} = 8\delta^{18}\text{O} + 10$ ) (Craig, 1961), and plotted the scale of isotopic changes from all sites in Figure 5. Interestingly, except for the 8.2 ka event, the isotopic excursions in New England appear to display the largest signal in the world, especially for the largest and longest 9.2 ka and 10.2 events when the isotopic changes in New England are several times greater than corresponding changes in Greenland, Asia and Europe. The isotopic signal of the 8.2 ka event in New England is modest compared with other records, which may be related to the different routing (i.e. Hudson Bay) of freshwater outburst at the final collapse of the LIS. The high amplitude of climate response to AMOC changes in eastern north America is consistent with model results (Shindell et al., 1999).

### Summary

The variability of decadal- to multidecadal-scale hydrogen isotope records from the northeastern USA coincides with key intervals of weakened AMOC, demonstrating the profound impact of AMOC oscillations on the abrupt continental climate changes in eastern North America. Our high resolution hydrogen isotopic records indicate that New England is highly sensitive to AMOC variations, and thus likely to experience significant climatic variations should the predicted major disturbance of AMOC occur in the coming century due to anthropogenic gas emissions.

### Acknowledgements

We thank Drs James Russell, Enno Schefuß and two reviewers for critical comments.



**Figure 5.** Compilation of selected isotope records showing the abrupt climate reversals during the early Holocene around the world, at (A) 8.2 ka, (B) 8.8 ka, (C) 9.2 ka, (D) 9.6 ka, (E) 10.2 ka and (F) 10.6 ka. Solid circles represent stable isotope records ( $\delta^{18}\text{O}$  and  $\delta\text{D}$ ). The size of the solid circles represents the amplitude of isotope reversal of individual records. Open circles represent non-isotope records

## Funding

This work was supported by grants from the National Science Foundation (NSF 0318050) to YH.

## References

- Alley RB and Agustsdottir AM (2005) The 8k event: Cause and consequences of a major Holocene abrupt climate change. *Quaternary Science Reviews* 24: 1123–1149.
- Bamberg A, Resenthal Y, Paul A, Heslop D, Mulitza S, Rühlemann C et al. (2010) Reduced North Atlantic Central Water formation in response to early Holocene ice-sheet melting. *Geophysical Research Letters* 37: L17705, doi:10.1029/2010GL043878.
- Bond G, Kromer B, Beer J, Muscheler R, Evans MN, Showers W et al. (2001) Persistent solar influence on north Atlantic climate during the Holocene. *Science* 294: 2130–2136.
- Bond G, Showers W, Cheseby M, Lotti R, Almasi P, deMenocal P et al. (1999) A pervasive millennial-scale cycle in North Atlantic Holocene and glacial climates. *Science* 278: 1257–1266.
- Bowen GJ and Revenaugh J (2003) Interpolating the isotopic composition of modern meteoric precipitation. *Water Resources Research* 39: doi:10.1029/2003WR002086.
- Clark PU, Marshall SJ, Clarke GKC, Hostetler SW, Licciardi JM and Teller JT (2001) Freshwater forcing of abrupt climate change during the last glaciation. *Science* 293: 283–287.
- Clark PU, Pisias NG, Stocker TF and Weaver AJ (2002) The role of the thermohaline circulation in abrupt climate change. *Nature* 415: 863–869.
- Clarke GKC, Bush ABG and Bush JWM (2009) Freshwater discharge, sediment transport, and modeled climate impacts of the final drainage of glacial Lake Agassiz. *Journal of Climate* 22: 2161–2180.
- Craig H (1961) Isotopic variations in meteoric waters. *Science* 133: 1702–1703.
- Daley T, Street-Perrott FA, Loader NJ, Barber KE, Hughes PDM, Fisher EH et al. (2009) Terrestrial climate signal of the ‘8200 yr B.P. cold event’ in the Labrador Sea region. *Geology* 37: 831–834.
- Dyke AS and Prest VK (1987) Late Wisconsinan and Holocene history of the Laurentide ice sheet. *Geographie physique et Quaternaire* 41: 237–263.
- Dykoski CA, Edwards RL, Cheng H, Yuan D, Cai Y, Zhang M et al. (2005) A high-resolution, absolute-dated Holocene and deglacial Asian monsoon record from Dongge Cave, China. *Earth and Planetary Science Letters* 233: 71–86, doi:10.1016/j.epsl.2005.1001.1036.
- Ellison CRW, Chapman MR and Hall IR (2006) Surface and deep ocean interactions during the cold climate event 8200 years ago. *Science* 312: 1929–1932.
- Faison EK, Foster DR, Oswald WW, Hansen BCS and Doughty E (2006) Early Holocene openlands in southern New England. *Ecology* 87: 2537–2547.
- Fleitmann D, Mudelsee M, Burns S, Bradley R, Kramers J and Matter A (2008) Evidence for a widespread climatic anomaly at around 9.2 ka before present. *Paleoceanography* 23: 6.
- Haeberli W and Gruber S (2009) Global warming and mountain permafrost. In: Margesin R (ed.) *Permafrost Soils*. Springer, 205–218.
- Hou J, Huang Y, Oswald WW, Foster DR and Shuman B (2007) Centennial-scale compound-specific hydrogen isotope record of Pleistocene–Holocene climate transition from southern New England. *Geophysical Research Letters* 34: doi:10.1029/2007GL030303.
- Hou J, Huang Y, Wang Y, Shuman B, Oswald WW, Faison E et al. (2006) Postglacial climate reconstruction based on compound-specific D/H ratios of fatty acids from Blood Pond, New England. *Geochemistry Geophysics Geosystems* 7: doi:10.1029/2005GC001076.
- Hu FS, Kaufma D, Yoneji S, Nelson D, Shemesh A, Huang Y et al. (2003) Cyclic variation and solar forcing of Holocene climate in the Alaskan subarctic. *Science* 301: 1890–1893.
- Huang Y, Shuman B, Wang Y and Webb T III (2002) Hydrogen isotope ratios of palmitic acid in lacustrine sediments record late Quaternary climate variations. *Geology* 30: 1103–1106.
- Hughen KA, Southon JR, Lehman SJ and Overpeck JT (2000) Synchronous radiocarbon and climate shifts during the last deglaciation. *Science* 290: 1951–1954.
- Kaser G, Hardy DR, Molg T, Bradley RS and Hayra TM (2004) Modern glacier retreat on Kilimanjaro as evidence of climate change: Observation and facts. *International Journal of Climatology* 24: 329–339.
- Kurek J, Cwynar LC and Spear RW (2004) The 8200 cal yr BP cooling event in eastern North America and the utility of midge analysis for Holocene temperature reconstructions. *Quaternary Science Reviews* 23: 627–639.
- LeGrande AN, Schmidt GA, Shindell DT, Field CV, Miller RL, Koch DM et al. (2006) Consistent simulations of multiple proxy responses to an abrupt climate change event. *Proceedings of National Academy of Sciences of the United States of America* 103: 837–842.
- Lutz B, Wiles G, Lowell T and Michaels J (2007) The 8.2 ka abrupt climate change event in Brown’s Lake, northeast Ohio. *Quaternary Research* 67: 292–296, doi:10.1016/j.yqres.2006.1008.1007.
- Namias J (1966) Nature and possible causes of the northeastern United States drought during 1962–65. *Monthly Weather Reviews* 94: 543–554.
- Newby P, Donnelly J, Shuman B and MacDonald D (2009) Evidence of centennial-scale drought from southeastern Massachusetts during the Pleistocene/Holocene transition. *Quaternary Science Reviews* 28: 1675–1692.
- Overpeck JT and Cole JE (2006) Abrupt change in Earth’s climate system. *Annual Review of Environment and Resources* 31: 1–31.
- Rhines P, Hakkinen S and Josey SA (2008) Is oceanic heat transport significant in the climate system. In: Dickson RR, Meincke J and Rhines P (eds) *Arctic-Subarctic Ocean Fluxes – Defining the Role of Northern Seas in Climate*. Springer, 87–109.
- Schmidt GA, Hoffmann G, Shindell DT and Hu YY (2005) Modeling atmospheric stable water isotopes and the potential for constraining cloud processes and stratosphere–troposphere water exchange. *Journal of Geophysical Research-Atmospheres* 110: D21314, doi:10.1029/2005JD005790.
- Seager R and Battisti DS (2007) Challenges to our understanding of the general circulation: Abrupt climate change. In: Schneider T and Sobel AS (eds) *The Global Circulation of the Atmosphere: Phenomena, Theory, Challenges*. Princeton University Press, 331–371.
- Severinghaus JP, Beaudette R, Headly MA, Taylor KC and Brook EJ (2009) Oxygen-18 of O<sub>2</sub> records the impact of abrupt climate change on the terrestrial biosphere. *Science* 324: 1431–1434.
- Shindell DT, Miller RL, Schmidt GA and Pandolfo L (1999) Simulation of recent northern winter climate trends by greenhouse-gas forcing. *Nature* 399: 452–455.
- Shuman B and Donnelly JP (2006) The influence of seasonal precipitation and temperature regimes on lake levels in the northeastern United States during the Holocene. *Quaternary Research* 65: 44–56.
- Shuman B, Bartlein PJ, Logar N, Newby P and Webb T III (2002) Parallel climate and vegetation responses to the early Holocene collapse of the Laurentide Ice Sheet. *Quaternary Science Reviews* 21: 1793–1805.
- Spurk M, Leuschner HH, Baillie MGL, Briffa KR and Friedrich M (2002) Depositional frequency of German subfossil oaks: Climatically and non-climatically induced fluctuations in the Holocene. *The Holocene* 12: 707–715.
- Stouffer RJ, Yin J, Gregory JM, Dixon KW, Spelman MJ, Hurlin W et al. (2006) Investigating the causes of the response of the Thermohaline Circulation to past and future climate changes. *Journal of Climate* 19: 1365–1387.
- Stuiver M, Reimer PJ, Bard E, Beck JW, Burr GS, Hughen KA et al. (1998) INTCAL98 radiocarbon age calibration, 24,000–0 cal BP. *Radiocarbon* 40: 1041–1083.
- Teller JT and Leverington DW (2004) Glacial Lake Agassiz: A 5000 yr history of change and its relationship to the  $\delta^{18}O$  record of Greenland. *Geological Society of America Bulletin* 116: 729.
- Vinther B, Clausen H, Johnsen S, Rasmussen S, Andersen K, Buchardt S et al. (2006) A synchronized dating of three Greenland ice cores throughout the Holocene. *Journal of Geophysical Research* 111: 11.
- von Grafenstein U, Erlenkeuser H, Brauer A, Jouzel J and Johnsen SJ (1999) A mid-European decadal isotope-climate record from 15,500 to 5000 years BP. *Science* 284: 1654–1657.
- Willemse NW and Tornqvist TE (1999) Holocene century-scale temperature variability from West Greenland lake records. *Geology* 27: 580–584.
- Xu W and Liu X (2007) Response of vegetation in the Qinghai-Tibet Plateau to global warming. *Chinese Geographical Sciences* 17: 151–159.
- Yancheva G, Nowaczyk N, Mingram J, Dulski P, Schettler G, Negendank J et al. (2007) Influence of the intertropical convergence zone on the East Asian monsoon. *Nature* 445: 74–77.
- Yu S-y, Colman SM, Lowell TV, Milne GA, Fisher TG, Breckenridge A et al. (2010) Freshwater outburst from Lake Superior as a trigger for the cold event 9300 years ago. *Science* 328: 1262–1266.