

West Tisbury

Produced in 2012

This report and associated map provide information about important sites for biodiversity conservation in your area.

This information is intended for conservation planning, and is <u>not</u> intended for use in state regulations.









Table of Contents

Introduction

What is *BioMap2* – Purpose and applications

One plan, two components

Understanding Core Habitat and its components

Understanding Critical Natural Landscape and its components

Understanding Core Habitat and Critical Natural Landscape Summaries

Sources of Additional Information

West Tisbury Overview


Core Habitat and Critical Natural Landscape Summaries

Elements of BioMap2 Cores

Core Habitat Summaries

Elements of *BioMap2* Critical Natural Landscapes

Critical Natural Landscape Summaries



Introduction

The Massachusetts Department of Fish & Game, through the Division of Fisheries and Wildlife's Natural Heritage & Endangered Species Program (NHESP), and The Nature Conservancy's Massachusetts Program developed *BioMap2* to protect the state's biodiversity in the context of climate change.


BioMap2 combines NHESP's 30 years of rigorously documented rare species and natural community data with spatial data identifying wildlife species and habitats that were the focus of the Division of Fisheries and Wildlife's 2005 State Wildlife Action Plan (SWAP). BioMap2 also integrates The Nature Conservancy's assessment of large, well-connected, and intact ecosystems and landscapes across the Commonwealth, incorporating concepts of ecosystem resilience to address anticipated climate change impacts.

Protection and stewardship of *BioMap2* Core Habitat and Critical Natural Landscape is essential to safeguard the diversity of species and their habitats, intact ecosystems, and resilient natural landscapes across Massachusetts.

What Does Status Mean?

The Division of Fisheries and Wildlife determines a status category for each rare species listed under the Massachusetts Endangered Species Act (MESA), M.G.L. c.131A, and its implementing regulations 321 CMR 10.00. Rare species are categorized as Endangered, Threatened or of Special Concern according to the following:


 Endangered species are in danger of extinction throughout all or a significant portion of their range or are in danger of extirpation from Massachusetts.



Get your copy of the *BioMap2* report! Download from www.nhesp.org or contact Natural Heritage at 508-389-6360 or natural.heritage@state.ma.us.

- Threatened species are likely to become Endangered in Massachusetts in the foreseeable future throughout all or a significant portion of their range.
- Special Concern species have suffered a
 decline that could threaten the species if
 allowed to continue unchecked or occur in
 such small numbers or with such restricted
 distribution or specialized habitat
 requirements that they could easily become
 Threatened in Massachusetts.

In addition NHESP maintains an unofficial watch list of plants that are tracked due to potential conservation interest or concern, but are <u>not</u> regulated under the Massachusetts Endangered Species Act or other laws or regulations. Likewise, described natural communities are <u>not</u> regulated by any law or regulations, but they can help to identify



Massachusetts Division of Fisheries and Wildlife

ecologically important areas that are worthy of protection. The status of natural communities reflects the documented number and acreages of each community type in the state:

- Critically Imperiled communities typically have 5 or fewer documented good sites or have very few remaining acres in the state.
- Imperiled communities typically have 6-20 good sites or few remaining acres in the state.
- Vulnerable communities typically have 21-100 good sites or limited acreage across the
- Secure communities typically have over 100 sites or abundant acreage across the state; however, excellent examples are identified as Core Habit to ensure continued protection.

In 2005 the Massachusetts Division of Fisheries and Wildlife completed a comprehensive State Wildlife Action Plan (SWAP) documenting the status of Massachusetts wildlife and providing recommendations to help guide wildlife conservation decision-making. SWAP includes all the wildlife species listed under the Massachusetts Endangered Species Act (MESA), as well as more than 80 species that need conservation attention but do not meet the requirements for inclusion under MESA. The SWAP document is organized around habitat types in need of conservation within the Commonwealth. While the original BioMap focused primarily on rare species protected under MESA, BioMap2 also addresses other Species of Conservation Concern, their habitats, and the ecosystems that support them to create a spatial representation of most of the elements of SWAP.

BioMap2: One Plan, Two Components

BioMap2 identifies two complementary spatial layers, Core Habitat and Critical Natural Landscape.

Core Habitat identifies key areas that are critical for the long-term persistence of rare species and other Species of Conservation Concern, as well as a wide diversity of natural communities and intact ecosystems across the Commonwealth. Protection of Core Habitats will contribute to the conservation of specific elements of biodiversity.

Critical Natural Landscape identifies large natural Landscape Blocks that are minimally impacted by development. If protected, these areas will provide habitat for wide-ranging native species, support intact ecological processes, maintain connectivity among habitats, and enhance ecological resilience to natural and anthropogenic disturbances in a rapidly changing world. Areas delineated as Critical Natural Landscape also include buffering upland around wetland, coastal, and aquatic Core Habitats to help ensure their longterm integrity.


The long-term persistence of Massachusetts biological resources requires a determined commitment to land and water conservation. Protection and stewardship of both Critical Natural Landscapes and Core Habitats are needed to realize the biodiversity conservation vision of BioMap2.

Components of Core Habitat

Core Habitat identifies specific areas necessary to promote the long-term persistence of rare species, other Species of Conservation Concern, exemplary natural communities, and intact ecosystems.

Rare Species

There are 432 native plant and animal species listed as Endangered, Threatened or Special Concern under the Massachusetts Endangered Species Act (MESA) based on their rarity, population trends, and threats to survival. For



Massachusetts Division of Fisheries and Wildlife



Table 1. Species of Conservation Concern described in the State Wildlife Action Plan and/or included on the MESA List and for which habitat was mapped in *BioMap2*. Note that plants are not included in SWAP, and that marine species such as whales and sea turtles are not included in *BioMap2*.

MESA-	Non-listed Species
listed	of Conservation
Species	Concern
4	5
27	23
10	5
4	3
10	17
102	9
256	0
413	62
	listed Species 4 27 10 4 10 102 256

BioMap2, NHESP staff identified the highest quality habitat sites for each non-marine species based on size, condition, and landscape context.

Other Species of Conservation Concern

In addition to species on the MESA List described previously, the State Wildlife Action Plan (SWAP) identifies 257 wildlife species and 22 natural habitats most in need of conservation within the Commonwealth. *BioMap2* includes species-specific habitat areas for 45 of these species and habitat for 17 additional species which was mapped with other coarse-filter and fine-filter approaches.

Priority Natural Communities

Natural communities are assemblages of plant and animal species that share a common environment and occur together repeatedly on the landscape. *BioMap2* gives conservation priority to natural communities with limited distribution and to the best examples of more common types.

Vernal Pools

Vernal pools are small, seasonal wetlands that provide important wildlife habitat, especially for amphibians and invertebrate animals that use them to breed. *BioMap*2 identifies the top 5 percent most interconnected clusters of Potential Vernal Pools in the state.

Forest Cores


In *BioMap2*, Core Habitat includes the best examples of large, intact forests that are least impacted by roads and development, providing critical habitat for numerous woodland species. For example, the interior forest habitat defined by Forest Cores supports many bird species sensitive to the impacts of roads and development, such as the Black-throated Green Warbler, and helps maintain ecological processes found only in unfragmented forest patches.

Wetland Cores

BioMap2 used an assessment of Ecological Integrity to identify the least disturbed wetlands in the state within undeveloped landscapes—those with intact buffers and little fragmentation or other stressors associated with development. These wetlands are most likely to support critical wetland functions (i.e., natural hydrologic conditions, diverse plant and animal habitats, etc.) and are most likely to maintain these functions into the future.

Aquatic Cores

To delineate integrated and functional ecosystems for fish species and other aquatic



Massachusetts Division of Fisheries and Wildlife

Species of Conservation Concern, beyond the species and exemplary habitats described above, *BioMap2* identifies intact river corridors within which important physical and ecological processes of the river or stream occur.

Components of Critical Natural Landscape

Critical Natural Landscape identifies intact landscapes in Massachusetts that are better able to support ecological processes and disturbance regimes, and a wide array of species and habitats over long time frames.

Landscape Blocks

BioMap2 identifies the most intact large areas of predominately natural vegetation, consisting of contiguous forests, wetlands, rivers, lakes, and ponds, as well as coastal habitats such as barrier beaches and salt marshes.

Upland Buffers of Wetland and Aquatic Cores

A variety of analyses were used to identify protective upland buffers around wetlands and rivers.

Upland Habitat to Support Coastal Adaptation

BioMap2 identifies undeveloped lands adjacent to and up to one and a half meters above existing salt marshes as Critical Natural Landscapes with high potential to support inland migration of salt marsh and other coastal habitats over the coming century.


The conservation areas identified by *BioMap2* are based on breadth and depth of data, scientific expertise, and understanding of Massachusetts' biodiversity. The numerous sources of information and analyses used to

Legal Protection of Biodiversity

BioMap2 presents a powerful vision of what Massachusetts would look like with full protection of the land most important for supporting the Commonwealth's biodiversity. While *BioMap2* is a planning tool with *no* regulatory function, all state-listed species enjoy legal protection under the Massachusetts Endangered Species Act (M.G.L. c.131A) and its implementing regulations (321 CMR 10.00). Wetland habitat of state-listed wildlife is also protected under the Wetlands Protection Act Regulations (310 CMR 10.00). The Natural Heritage Atlas contains maps of Priority Habitats and Estimated Habitats, which are used, respectively, for regulation under the Massachusetts Endangered Species Act and the Wetlands Protection Act. For more information on rare species regulations, and to view Priority and Estimated Habitat maps, please see the Regulatory Review page at www.mass.gov/dfwele/dfw/nhesp/regulatory_rev iew/reg review home.htm.

BioMap2 is a conservation planning tool that does not, in any way, supplant the Estimated and Priority Habitat Maps which have regulatory significance. Unless and until the BioMap2 vision is fully realized, we must continue to protect our most imperiled species and their habitats.

create Core Habitat and Critical Natural
Landscape are complementary, and outline a
comprehensive conservation vision for
Massachusetts, from rare species to intact
landscapes. In total, these robust analyses define
a suite of priority lands and waters that, if
permanently protected, will support
Massachusetts' natural systems for generations
to come.



Massachusetts Division of Fisheries and Wildlife

Understanding Core Habitat Summaries

Following the Town Overview, there is a descriptive summary of each Core Habitat and Critical Natural Landscape that occurs in your city or town. These summaries highlight some of the outstanding characteristics of each Core Habitat and Critical Natural Landscape, and will help you learn more about your city or town's biodiversity. You can find out more information about many of these species and natural communities by looking at specific fact sheets at www.nhesp.org.

Additional Information

For copies of the full *BioMap2* report, the Technical Report, and an <u>interactive mapping tool</u>, visit the *BioMap2* <u>website</u> via the Land Protection and Planning tab at <u>www.nhesp.org</u>. If you have any questions about this report, or if you need help protecting land for biodiversity in your community, the Natural Heritage & Endangered Species Program staff looks forward to working with you.

Contact the Natural Heritage & Endangered Species Program

By phone 508-389-6360 By fax 508-389-7890

By email <u>natural.heritage@state.ma.us</u>
By Mail 100 Hartwell Street, Suite 230
West Boylston, MA 01583


The GIS datalayers of *BioMap2* are available for download from MassGIS at www.mass.gov/mgis.



phone: 508-389-6360 fax: 508-389-7890

Town Overview

West Tisbury lies within the Cape Cod and Islands Ecoregion, an area formed by three advances and retreats of the Wisconsin Ice Sheet. The resulting terminal moraines, outwash plains, and coastal deposits characterize the area with their sandy beaches, grassy dunes, bays, marshes, and scrubby oak-pine forests. There are numerous kettle hole ponds, swamps, and bogs. Much of the surface water is highly acidic.



West Tisbury at a Glance

- Total Area: 16,823 acres (26.3 square miles)
- Human Population in 2010: 2,740
- Open space protected in perpetuity: 6,576 acres, or 39.1% percent of total area*
- BioMap2 Core Habitat: 12,938 acres
- BioMap2 Core Habitat Protected: 6,131 acres or 47.4%
- *BioMap2* Critical Natural Landscape: 10,853 acres
- *BioMap2* Critical Natural Landscape Protected: 6,102 acres or 56.2%.


BioMap2 Components

Core Habitat

- 3 Exemplary or Priority Natural Community Cores
- 1 Forest Core
- 7Aquatic Cores
- 7 Species of Conservation Concern Cores**
 - 8 birds, 3 reptiles, 1 amphibian, 1 fish, 24 insects, 18 plants


Critical Natural Landscape

- 4 Landscape Blocks
- 6 Aquatic Core Buffers
- 2 Coastal Adaptation Areas
- 8Tern Foraging Areas
- * Calculated using MassGIS data layer "Protected and Recreational Open Space—March, 2012".
- ** See next pages for complete list of species, natural communities and other biodiversity elements.





BioMap2 Core Habitat and Critical Natural Landscape in West Tisbury






BioMap2 Core Habitat



BioMap2 Critical Natural Landscape

1 Mile





Species of Conservation Concern, Priority and Exemplary Natural Communities, and Other Elements of Biodiversity in West Tisbury

Insects

Moths

Coastal Heathland Cutworm, (Abagrotis nefascia), SC

Barrens Daggermoth, (Acronicta albarufa), T

Gerhard's Underwing, (Catocala herodias gerhardi), SC

Melsheimer's Sack Bearer, (Cicinnus melsheimeri), T

Chain Dot Geometer, (Cingilia catenaria), SC

Imperial Moth, (Eacles imperialis), T

Slender Clearwing Sphinx Moth, (Hemaris gracilis), SC

Barrens Buckmoth, (Hemileuca maia), SC

Coastal Swamp Metarranthis, (Metarranthis pilosaria), SC

Water-willow Stem Borer, (Papaipema sulphurata), T

Spartina Borer, (Photedes inops), SC

Pink Sallow Moth, (Psectraglaea carnosa), SC

Pine Barrens Zale, (Zale lunifera), SC

Barrens Metarranthis, (Metarranthis apiciaria), E

Dune Noctuid Moth, (Sympistis riparia), SC

Faded Gray Geometer, (Stenoporpia polygrammaria), T

Pine Barrens Lycia, (Lycia ypsilon), T

Sandplain Euchlaena, (Euchlaena madusaria), SC

Sandplain Heterocampa, (Heterocampa varia), T

Southern Ptichodis, (Ptichodis bistrigata), T

Three-lined Angle Moth, (Digrammia eremiata), T

Unexpected Cycnia, (Cycnia inopinatus), T

Beetles

Purple Tiger Beetle, (Cicindela purpurea), SC

Dragonflies

Comet Darner, (Anax longipes), SC

Amphibians

Four-toed Salamander, (Hemidactylium scutatum), Non-listed SWAP

Fishes


American Brook Lamprey, (Lampetra appendix), T

Reptiles

Eastern Box Turtle, (Terrapene carolina), SC

Smooth Green Snake, (Opheodrys vernalis), Non-listed SWAP

Spotted Turtle, (Clemmys guttata), Non-listed SWAP



Massachusetts Division of Fisheries and Wildlife

Birds

Grasshopper Sparrow, (Ammodramus savannarum), T

Piping Plover, (Charadrius melodus), T

Northern Harrier, (Circus cyaneus), T

Common Tern, (Sterna hirundo), SC

Roseate Tern, (Sterna dougallii), E

Least Tern, (Sternula antillarum), SC

Barn Owl, (Tyto alba), SC

Eastern Whip-poor-will, (Caprimulgus vociferus), SC

Plants

Hairy Agrimony, (Agrimonia pubescens), T

Purple Needlegrass, (Aristida purpurascens), T

Pygmyweed, (Crassula aquatica), T

Bushy Rockrose, (Crocanthemum dumosum), SC

Saltpond Pennywort, (Hydrocotyle verticillata), T

Saltpond Grass, (Leptochloa fusca ssp. fascicularis), T

Sandplain Flax, (Linum intercursum), SC

Bayard's Green Adder's-mouth, (Malaxis bayardii), E

Canadian Sanicle, (Sanicula canadensis), T

Papillose Nut-sedge, (Scleria pauciflora), E

Bristly Foxtail, (Setaria parviflora), SC

Sandplain Blue-eyed Grass, (Sisyrinchium fuscatum), SC

Grass-leaved Ladies'-tresses, (Spiranthes vernalis), T

Cranefly Orchid, (Tipularia discolor), E

Lion's Foot, (Nabalus serpentarius), E

Nantucket Shadbush, (Amelanchier nantucketensis), recently de-listed

River Arrowhead, (Sagittaria subulata), recently de-listed

Sea-beach Knotweed, (Polygonum glaucum), SC

Priority Natural Communities

Estuarine Subtidal: Coastal Salt Pond, S2

Other BioMap2 Components

Forest Core

Aquatic Core

Landscape Block

Aquatic Core Buffer

Coastal Adaptation Area

Tern Foraging Area

E = Endangered




Massachusetts Division of Fisheries and Wildlife




Conserving the Biodiversity of Massachusetts in a Changing World


- T = Threatened
- SC = Special Concern
- S1 = Critically Imperiled communities, typically 5 or fewer documented sites or very few remaining acres in the state.
- S2 = Imperiled communities, typically 6-20 sites or few remaining acres in the state.
- S3 = Vulnerable communities, typically have 21-100 sites or limited acreage across the state.



BioMap2 Core Habitat in West Tisbury

Core IDs correspond with the following element lists and summaries.





1 Mile





Elements of BioMap2 Cores


This section lists all elements of *BioMap2* Cores that fall *entirely or partially* within West Tisbury. The elements listed here may not occur within the bounds of West Tisbury.

Core 56


	Species of Conservation Concern Water-willow Stem Borer	Papaipema sulphurata	T
Core 8	32		
	Species of Conservation Concern Faded Gray Geometer Gerhard's Underwing Moth	Stenoporpia polygrammaria Catocala herodias gerhardi	T SC
Core 8	66		
	Species of Conservation Concern Unexpected Cycnia	Cycnia inopinatus	T
Core 8	37		
	Species of Conservation Concern Faded Gray Geometer Gerhard's Underwing Moth Water-willow Stem Borer	Stenoporpia polygrammaria Catocala herodias gerhardi Papaipema sulphurata	T SC T
Core 8	8		
	Species of Conservation Concern Barrens Daggermoth Faded Gray Geometer Gerhard's Underwing Moth Pine Barrens Zale	Acronicta albarufa Stenoporpia polygrammaria Catocala herodias gerhardi Zale lunifera	T T SC SC
Core 8	89		
	Aquatic Core Species of Conservation Concern Saltpond Grass Least Tern	Leptochloa fusca ssp. fascicularis Sternula antillarum	T SC
Cara 1	03		

Core 102


Forest Core Wetland Core Aquatic Core



Massachusetts Division of Fisheries and Wildlife




Priority & Exemplary Natural Communi	ities	
Estuarine Subtidal: Coastal Salt Pond	d	S2
Sandplain Grassland		S1
Species of Conservation Concern		
Bayard's Green Adder's-mouth	Malaxis bayardii	E
Bristly Foxtail	Setaria parviflora	SC
Bushy Rockrose	Crocanthemum dumosum	SC
Canadian Sanicle	Sanicula canadensis	T
Cranefly Orchid	Tipularia discolor	E
Grass-leaved Ladies'-tresses	Spiranthes vernalis	T
Hairy Agrimony	Agrimonia pubescens	T
Lion's Foot	Nabalus serpentarius	E
New England Blazing Star	Liatris scariosa var. novae-angliae	SC
Northern Gama-grass	Tripsacum dactyloides	E
Papillose Nut Sedge	Scleria pauciflora	E
Purple Needlegrass	Aristida purpurascens	T
Pygmyweed	Crassula aquatica	T
Saltpond Pennywort	Hydrocotyle verticillata	T
Sandplain Blue-eyed Grass	Sisyrinchium fuscatum	SC
Sandplain Flax	Linum intercursum	SC
Sea-beach Knotweed	Polygonum glaucum	SC
Smooth Branched Sponge	Spongilla aspinosa	SC
Barrens Buckmoth	Hemileuca maia	SC
Barrens Daggermoth	Acronicta albarufa	T
Barrens Metarranthis Moth	Metarranthis apiciaria	E
Chain Dot Geometer	Cingilia catenaria	SC
Coastal Heathland Cutworm	Abagrotis nefascia	SC
Coastal Swamp Metarranthis Moth	Metarranthis pilosaria	SC
Dune Noctuid Moth	Sympistis riparia	SC
Faded Gray Geometer	Stenoporpia polygrammaria	T
Gerhard's Underwing Moth	Catocala herodias gerhardi	SC
Imperial Moth	Eacles imperialis	T
Melsheimer's Sack Bearer	Cicinnus melsheimeri	T
Pine Barrens Lycia	Lycia ypsilon	T
Pine Barrens Speranza	Speranza exonerata	SC
Pine Barrens Zale	Zale lunifera	SC
Pink Sallow	Psectraglaea carnosa	SC
Sandplain Euchlaena	Euchlaena madusaria	SC
Sandplain Heterocampa	Heterocampa varia	Non-listed SWAP
Slender Clearwing Sphinx Moth	Hemaris gracilis	SC
Southern Ptichodis	Ptichodis bistrigata	T
Spartina Borer Moth	Photedes inops	SC
The Pink Streak	Dargida rubripennis	T
Three-lined Angle Moth	Digrammia eremiata	T



Massachusetts Division of Fisheries and Wildlife



Unexpected Cycnia	Cycnia inopinatus	T
Water-willow Stem Borer	Papaipema sulphurata	T
Oak Hairstreak	Satyrium favonius	SC
Purple Tiger Beetle	Cicindela purpurea	SC
Comet Darner	Anax longipes	SC
Eastern Spadefoot	Scaphiopus holbrookii	T
Four-toed Salamander	Hemidactylium scutatum	Non-listed SWAP
Eastern Box Turtle	Terrapene carolina	SC
Eastern Ribbon Snake	Thamnophis sauritus	Non-listed SWAP
Smooth Green Snake	Opheodrys vernalis	Non-listed SWAP
Spotted Turtle	Clemmys guttata	Non-listed SWAP
American Brook Lamprey	Lampetra appendix	T
Barn Owl	Tyto alba	SC
Common Tern	Sterna hirundo	SC
Eastern Whip-poor-will	Caprimulgus vociferus	SC
Grasshopper Sparrow	Ammodramus savannarum	T
Least Tern	Sternula antillarum	SC
Northern Harrier	Circus cyaneus	T
Piping Plover	Charadrius melodus	T
Roseate Tern	Sterna dougallii	E



Core Habitat Summaries

Core 56

A 3-acre Core Habitat featuring a Species of Conservation Concern.

The Water-willow Stem Borer is a yellowish moth with purple-brown shading that inhabits shallow portions of coastal plain wetlands where water-willow grows. It is endemic to southeastern Massachusetts.

Core 82

A 55-acre Core Habitat featuring Species of Conservation Concern.

On Martha's Vineyard, the Faded Gray Geometer inhabits scrub oak barrens and open oak woodland. Larval hosts are undocumented in Massachusetts, but are probably oaks (Quercus spp.).

Gerhard's Underwing, a noctuid moth, inhabits xeric and open pitch pine/scrub oak barrens, especially scrub oak thickets on sandplains or rocky summits and ridges. The larvae feed on the catkins and new leaves of scrub oak (Quercus ilicifolia), and must complete feeding in the spring before the catkins drop off and the new leaves harden.

Core 86

A 11-acre Core Habitat featuring a Species of Conservation Concern.

The Unexpected Cycnia inhabits open, grassy habitats with abundant milkweeds (Asclepias spp.), especially sandplain grasslands. Larvae feed on milkweeds (Asclepias spp.), especially butterfly weed (A. tuberosa).


Core 87

A 62-acre Core Habitat featuring Species of Conservation Concern.

On Martha's Vineyard, the Faded Gray Geometer inhabits scrub oak barrens and open oak woodland. Larval hosts are undocumented in Massachusetts, but are probably oaks (Quercus spp.).

Gerhard's Underwing, a noctuid moth, inhabits xeric and open pitch pine/scrub oak barrens, especially scrub oak thickets on sandplains or rocky summits and ridges. The larvae feed on the catkins and new leaves of scrub oak (Quercus ilicifolia), and must complete feeding in the spring before the catkins drop off and the new leaves harden.

The Water-willow Stem Borer is a yellowish moth with purple-brown shading that inhabits shallow portions of coastal plain wetlands where water-willow grows. It is endemic to southeastern Massachusetts.



Massachusetts Division of Fisheries and Wildlife 100 Hartwell Street, Suite 230, West Boylston, MA 01583

phone: 508-389-6360 fax: 508-389-7890

Core 88

A 20-acre Core Habitat featuring Species of Conservation Concern.

The Barrens Daggermoth, a noctuid moth, inhabits xeric, oak-dominated woodland, barrens, and scrub habitats on sandy soil.

On Martha's Vineyard, the Faded Gray Geometer inhabits scrub oak barrens and open oak woodland. Larval hosts are undocumented in Massachusetts, but are probably oaks (*Quercus* spp.).

Gerhard's Underwing, a noctuid moth, inhabits xeric and open pitch pine/scrub oak barrens, especially scrub oak thickets on sandplains or rocky summits and ridges. The larvae feed on the catkins and new leaves of scrub oak (*Quercus ilicifolia*), and must complete feeding in the spring before the catkins drop off and the new leaves harden.

The Pine Barrens Zale is found in sandplain pitch pine/scrub oak barrens, especially in scrub oak thickets. Larvae feed on scrub oak (*Quercus ilicifolia*).

Core 89

A 37-acre Core Habitat featuring Aquatic Core and Species of Conservation Concern.

Aquatic Cores are intact river corridors within which important physical and ecological processes of the river or stream occur. They delineate integrated and functional ecosystems for fish species and other aquatic Species of Conservation Concern.


In Massachusetts, Saltpond Grass is native to sandy shores of brackish coastal ponds, often where freshwater enters the pond such as at a seep or small stream.

Diminutive yet feisty, the Least Tern is a spring and summer colonial nester on Massachusetts' sandy beaches. For nesting, it favors for sites with little or no vegetation. In Massachusetts, the Least Tern nests on sandy or gravelly beaches periodically scoured by storm tides, resulting in sparse or no vegetation; it also takes advantage of dredge spoils. Along the coast, the Least Tern forages in shallow-water habitats, including bays, lagoons, estuaries, river and creek mouths, tidal marshes, and ponds.

Core 102

A 35,587-acre Core Habitat featuring Forest Core, Wetland Core, Aquatic Core, Priority Natural Communities, and Species of Conservation Concern.

Much of Martha's Vineyard is a large and complex Core Habitat, home to 65 rare and uncommon species, with 24 of those globally rare. In fact, the highest concentration of rare species in the state is near the center of the island in Corellus State Forest. Many of these rare species are moths, tiger beetles, and plants which inhabit only the kinds of sandplains that cover much of the island. As well, the Vineyard supports some of the best populations in the state of breeding Northern Harriers and Whip-poor-wills. The sandplains and barrens of this island have remained in states of early succession over the centuries



because of fire and major storms; nowadays, active management techniques, such as prescribed fire and mowing, help keep these singular habitats alive.


Coastal Salt Pond communities consist of vegetation surrounding, and in, coastal brackish ponds. These ponds are usually separated from the ocean by a sandspit. Their salinity varies and is influenced by opening and closing of the spit. Five examples of Coastal Salt Pond, mostly large, in excellent condition, and well buffered in naturally vegetated settings.

Sandplain Grasslands are open, essentially treeless, grass-dominated communities that generally occur on sand or other dry, poor soils. Occurrences are maintained by fire, salt spray, and, now, mowing. This Core has two examples of Sandplain Grassland including one that is considered to be the largest and best in the state, and possibly in New England. This very rare natural community harbors many state-listed plant and animal species.

Forest Cores are the best examples of large, intact forests that are least impacted by roads and development. Forest Cores support many bird species sensitive to the impacts of roads and development and help maintain ecological processes found only in unfragmented forest patches.


Wetland Cores are the least disturbed wetlands in the state within undeveloped landscapes—those with intact buffers and little fragmentation or other stressors associated with development. These wetlands are most likely to support critical wetland functions (i.e., natural hydrologic conditions, diverse plant and animal habitats, etc.) and are most likely to maintain these functions into the future.


Aquatic Cores are intact river corridors within which important physical and ecological processes of the river or stream occur. They delineate integrated and functional ecosystems for fish species and other aquatic Species of Conservation Concern.



BioMap2 Critical Natural Landscape in West Tisbury


Critical Natural Landscape IDs correspond with the following element lists and summaries.











Elements of *BioMap2* Critical Natural Landscapes

This section lists all elements of *BioMap2* Critical Natural Landscapes that fall *entirely or partially* within West Tisbury. The elements listed here may not occur within the bounds of West Tisbury.

CNL 40


Landscape Block

CNL 42

Landscape Block

CNL 45

Aquatic Core Buffer Coastal Adaptation Area Landscape Block Tern Foraging Area



Critical Natural Landscape Summaries

CNL 40

A 2,926-acre Critical Natural Landscape featuring Landscape Block.

Landscape Blocks, the primary component of Critical Natural Landscapes, are large areas of intact predominately natural vegetation, consisting of contiguous forests, wetlands, rivers, lakes, and ponds, as well as coastal habitats such as barrier beaches and salt marshes. Pastures and power-line rights-of-way, which are less intensively altered than most developed areas, were also included since they provide habitat and connectivity for many species. Collectively, these natural cover types total 3.6 million acres across the state. An Ecological Integrity assessment was used to identify the most intact and least fragmented areas. These large Landscape Blocks are most likely to maintain dynamic ecological processes such as buffering, connectivity, natural disturbance, and hydrological regimes, all of which help to support wide-ranging wildlife species and many other elements of biodiversity.


In order to identify critical Landscape Blocks in each ecoregion, different Ecological Integrity thresholds were used to select the largest intact landscape patches in each ecoregion while avoiding altered habitat as much as possible. This ecoregional representation accomplishes a key goal of *BioMap2* to protect the ecological stages that support a broad suite of biodiversity in the context of climate change. Blocks were defined by major roads, and minimum size thresholds differed among ecoregions to ensure that *BioMap2* includes the best of the best in each ecoregion.

CNL 42

A 534-acre Critical Natural Landscape featuring Landscape Block.

Landscape Blocks, the primary component of Critical Natural Landscapes, are large areas of intact predominately natural vegetation, consisting of contiguous forests, wetlands, rivers, lakes, and ponds, as well as coastal habitats such as barrier beaches and salt marshes. Pastures and power-line rights-of-way, which are less intensively altered than most developed areas, were also included since they provide habitat and connectivity for many species. Collectively, these natural cover types total 3.6 million acres across the state. An Ecological Integrity assessment was used to identify the most intact and least fragmented areas. These large Landscape Blocks are most likely to maintain dynamic ecological processes such as buffering, connectivity, natural disturbance, and hydrological regimes, all of which help to support wide-ranging wildlife species and many other elements of biodiversity.

In order to identify critical Landscape Blocks in each ecoregion, different Ecological Integrity thresholds were used to select the largest intact landscape patches in each ecoregion while avoiding altered habitat as much as possible. This ecoregional representation accomplishes a key goal of *BioMap2* to protect the ecological stages that support a broad suite of biodiversity in the context of climate change. Blocks were defined by major roads, and minimum size thresholds differed among ecoregions to ensure that *BioMap2* includes the best of the best in each ecoregion.



CNL 45

A 42,510-acre Critical Natural Landscape featuring Aquatic Core Buffer, Landscape Block, Coastal Adaptation Area, and Tern Foraging Area.

A variety of analyses were used to identify protective upland buffers around wetlands and rivers. One, the variable width buffers methodology, included the most intact areas around each wetland and river, by extending deeper into surrounding unfragmented habitats than into developed areas adjacent to each wetland. Other upland buffers were identified through the rare species habitat analysis. In this way, the conservation of wetland buffers will support the habitats and functionality of each wetland, and also include adjacent uplands that are important for many species that move between habitat types.


Landscape Blocks, the primary component of Critical Natural Landscapes, are large areas of intact predominately natural vegetation, consisting of contiguous forests, wetlands, rivers, lakes, and ponds, as well as coastal habitats such as barrier beaches and salt marshes. Pastures and power-line rights-of-way, which are less intensively altered than most developed areas, were also included since they provide habitat and connectivity for many species. Collectively, these natural cover types total 3.6 million acres across the state. An Ecological Integrity assessment was used to identify the most intact and least fragmented areas. These large Landscape Blocks are most likely to maintain dynamic ecological processes such as buffering, connectivity, natural disturbance, and hydrological regimes, all of which help to support wide-ranging wildlife species and many other elements of biodiversity.

In order to identify critical Landscape Blocks in each ecoregion, different Ecological Integrity thresholds were used to select the largest intact landscape patches in each ecoregion while avoiding altered habitat as much as possible. This ecoregional representation accomplishes a key goal of *BioMap2* to protect the ecological stages that support a broad suite of biodiversity in the context of climate change. Blocks were defined by major roads, and minimum size thresholds differed among ecoregions to ensure that *BioMap2* includes the best of the best in each ecoregion.

At 10,411 acres, this Landscape Block is the fourth largest in the ecoregion, supports critical habitat, and is especially important in the developed landscapes of the Massachusetts Islands.

The coastal habitats of Massachusetts are particularly vulnerable to potential sea-level rise in the next century, which many estimates suggest is likely to exceed one meter. Therefore, in addition to prioritizing current coastal habitats, the creators of *BioMap2* examined the landward side of salt marshes to determine where these habitats might move to as sea levels rise. Undeveloped lands adjacent to and up to one and a half meters above existing salt marshes were identified, and included as Critical Natural Landscapes with high potential to support inland migration of salt marsh and other coastal habitats over the coming century.

Terns range widely from their breeding colonies to forage. While the breeding and staging areas for Roseate, Arctic, Common, and Least Terns were included in the Species of Conservation Concern Core Habitat for *BioMap2*, tern foraging areas were included in *BioMap2* as part of Critical Natural Landscape. The extent of foraging habitat for Arctic, Common, and Roseate Terns depends on the size of the breeding colony. For Least Tern, all shallow marine and estuarine waters within 2 miles of recent colony sites and up to 1 mile offshore were mapped as foraging habitat.



Help Save Endangered Wildlife!

Please contribute on your Massachusetts income tax form or directly to the



Natural Heritage & Endangered Species Fund

To learn more about the Natural Heritage & Endangered Species Program and the Commonwealth's rare species, visit our web site at www.nhesp.org.