

Research



Cite this article: Park DS, Breckheimer I, Williams AC, Law E, Ellison AM, Davis CC. 2018 Herbarium specimens reveal substantial and unexpected variation in phenological sensitivity across the eastern United States. *Phil. Trans. R. Soc. B* **374**: 20170394. <http://dx.doi.org/10.1098/rstb.2017.0394>

Accepted: 15 October 2018

One contribution of 16 to a theme issue 'Biological collections for understanding biodiversity in the Anthropocene'.

Subject Areas:

ecology, evolution, taxonomy and systematics

Keywords:

citizen science, digitization, geographical range, herbarium specimens, phenology, phenological sensitivity

Authors for correspondence:

Daniel S. Park

e-mail: danielpark@fas.harvard.edu

Charles C. Davis

e-mail: cdavis@oeb.harvard.edu

[†]These authors contributed equally to this work.

Electronic supplementary material is available online at <https://dx.doi.org/10.6084/m9.figshare.c.4274660>.

Herbarium specimens reveal substantial and unexpected variation in phenological sensitivity across the eastern United States

Daniel S. Park^{1,†}, Ian Breckheimer^{1,†}, Alex C. Williams², Edith Law², Aaron M. Ellison³ and Charles C. Davis¹

¹Department of Organismic and Evolutionary Biology and Harvard University Herbaria, Harvard University, Cambridge, MA 02138, USA

²David R. Cheriton School of Computer Science, University of Waterloo, Waterloo, Ontario, Canada N2L 3G1

³Harvard Forest, Harvard University, Petersham, MA 01366, USA

CCD, 0000-0001-8747-1101

Phenology is a key biological trait that can determine an organism's survival and provides one of the clearest indicators of the effects of recent climatic change. Long time-series observations of plant phenology collected at continental scales could clarify latitudinal and regional patterns of plant responses and illuminate drivers of that variation, but few such datasets exist. Here, we use the web tool *CrowdCurio* to crowdsource phenological data from over 7000 herbarium specimens representing 30 diverse flowering plant species distributed across the eastern United States. Our results, spanning 120 years and generated from over 2000 crowdsourcers, illustrate numerous aspects of continental-scale plant reproductive phenology. First, they support prior studies that found plant reproductive phenology significantly advances in response to warming, especially for early-flowering species. Second, they reveal that fruiting in populations from warmer, lower latitudes is significantly more phenologically sensitive to temperature than that for populations from colder, higher-latitude regions. Last, we found that variation in phenological sensitivities to climate within species between regions was of similar magnitude to variation between species. Overall, our results suggest that phenological responses to anthropogenic climate change will be heterogeneous within communities and across regions, with large amounts of regional variability driven by local adaptation, phenotypic plasticity and differences in species assemblages. As millions of imaged herbarium specimens become available online, they will play an increasingly critical role in revealing large-scale patterns within assemblages and across continents that ultimately can improve forecasts of the impacts of climatic change on the structure and function of ecosystems.

This article is part of the theme issue 'Biological collections for understanding biodiversity in the Anthropocene'.

1. Introduction

Ecosystems on every continent have been affected by local, regional and global changes in climate, especially increases in temperature [1]. Changes in phenology—the timing of life-history events—are among the most conspicuous and well-documented species responses to climatic change, especially for plants [2–7]. Phenological disruption has already impacted species' local persistence and community diversity [8–10], which may have widespread consequences for critical ecosystem processes, including carbon sequestration [11–13], ecosystem–atmosphere interactions [14] and trophic interactions [15–28].

Despite these trends, our knowledge of plant phenological responses to climatic change remains inadequate. In particular, although phenological responses may differ among species with different functional or life-history traits and biogeographical origins [29–32], long-term observational datasets

to assess such trends are limited in geographical, temporal and taxonomic scope [33]. Many of these data track woody plant species of the Northern Hemisphere (most commonly, abundant tree species), and only for the last approximately 40 years (but see [34,35]). These biases limit our understanding of variation in phenological responses across species and biomes. Furthermore, although population-level variation in phenology has been demonstrated for a few species [36–38], there are very few studies that quantify both inter- and intraspecific variation in phenological response [32,33].

Variability in species' phenology is particularly relevant because climatic change is not geographically uniform. For example, high-latitude regions are warming faster than subtropical and tropical ones [1,39]. Short growing seasons also may cause high-latitude ecosystems to be especially sensitive to temperature, leading to stronger selective pressures for populations to initiate growth as soon as conditions become favourable in early spring. Additionally, plants adapted to highly variable climates may exhibit higher phenological thresholds to temperature, as it provides a less reliable signal [40]. Thus, the effects of climatic change on species' phenology may differ across their ranges depending on variation in phenological sensitivity. Variability in phenological responses to climatic change within species may also alter patterns of gene flow, which could either promote or counteract adaptive evolution via the sharing of locally (mal)adapted alleles [41–44]. Recent studies have shown that plant phenology may be more responsive in more northern-ranging populations where there are more variable and extreme climates, especially during the early part of the growing season [44,45]. Although these studies are suggestive, they are restricted in spatial scale and taxonomic scope, and broad regional patterns of phenological response to climate may differ from patterns at smaller, local scales.

Herbarium specimens represent snapshots of phenology (i.e. flowering and fruiting) at a specific place and time, and have shown tremendous promise to increase the spatial, temporal and taxonomic resolution of phenological data [46–48]. They provide rich historical depth, wide geographical scope and taxonomic diversity, all of which allow researchers to track long-term changes of vast numbers of species and communities through space and time [4,46–49]. Despite their representation of phenological responses [48,50], herbarium specimens have been used less frequently than other data sources, such as field observations, to address phenological change, in part because they have been inaccessible to many researchers [46,51]. However, the widespread digitization of herbarium collections [52] combined with new approaches to collecting [46,53] and analysing [54,55] phenological data derived from herbarium specimens has the power to transform our understanding of plant responses to global climatic change.

Here, we applied a newly developed web-enabled crowdsourcing platform, *CrowdCurio: Thoreau's Field Notes* (<https://www.crowdcurio.com/>) [46], to examine more than 7000 specimens of 30 phylogenetically diverse flowering plant species. The *Thoreau's Field Notes* module facilitates the rapid quantification of phenological traits via image annotation and has been demonstrated to yield reliable data regardless of the level of expertise among crowdsourcers (i.e. expert versus non-expert scoring) [46]. We used these crowdsourced data to infer the magnitude, direction, and variability in reproductive phenological responses to spring

temperature across 23° of latitude in the eastern United States. We examined both native and introduced plant species from northern coniferous forests, eastern deciduous forests, subtropical evergreen forests, grasslands, wetlands, alpine meadows and aquatic plant communities. Environmental conditions in this region vary considerably across species' ranges, and populations may exhibit substantial variation in phenological response across this latitudinal gradient. Our overall goals with this study were: (i) to demonstrate the power of characterizing phenology from herbarium data using an efficient and rapid workflow that leverages a nearly fully mobilized online flora of the eastern United States [56,57]; (ii) to greatly increase the taxonomic and ecological diversity of species sampled for this purpose (from woody perennials to herbaceous annuals across a range of biomes); and (iii) to sample species with broad latitudinal ranges to assess regional and inter- and intra-species variation in phenological responses.

2. Methods

(a) Specimen data collection

We examined phenological responses of species using digitized specimens from two of the most comprehensive digitized regional floras in the world, the Consortium of Northeastern Herbaria (CNH; <http://portal.neherbaria.org/portal/>) [56] and Southeast Regional Network of Expertise and Collections (SERNEC; <http://sernecportal.org/portal/index.php>) [57]. These two online portals include more than six million digitized herbarium records, including specimen images. Our criteria for selecting angiosperm species for analysis were that specimens: (i) included at least county-level location data; (ii) included at least 50 unique collections across space and time; (iii) were of species with relatively easily identifiable and quantifiable reproductive structures; and (iv) were from species with broad latitudinal ranges sufficient to enable quantification of population-level variation.

Applying these criteria yielded 30 species with varying life-history traits, growth forms, native status and general reproductive seasonality (e.g. early- versus late-spring flowering). We downloaded over 10 000 digital herbarium specimen images of these species from CNH and SERNEC, removed duplicate, misidentified or sterile specimens, and those with notable insect damage on reproductive structures, extensive physical damage or poor preservation. We also removed all 30 specimens from Florida, which were geographical and climatic outliers. Our final dataset comprised 7722 specimens and spanned 120 years across 512 United States counties (table 1). Species' life-history (annual versus perennial), growth form (woody versus herbaceous) and native status (native versus introduced) were inferred from the United States Department of Agriculture PLANTS Database (<https://plants.usda.gov/>). For individual specimen metadata, see electronic supplementary material, table S1.

(b) Crowdsourcing phenological data collection

The phenological state (phenophase) of a plant can be inferred from the presence and quantity of relevant structures, such as leaves, flowers or fruits [46,48]. Past researchers generally have focused on the presence or the absence of a single structure or trait (e.g. [58]) or applied majority estimates for scoring a single phenophase (e.g. [49]). Here, we quantified data for two reproductive phenophases, flowering and fruiting. Specimens were scored as flowering if open flowers represented greater than or equal to 50% of the total reproductive structure count

Table 1. The number of specimens and different categorical traits of examined species. FL and FR refer to the number of specimens classified as flowering and fruiting, respectively.

family	species	time span (years)	FL	FR	lifespan	growth form	status
Ranunculaceae	<i>Anemone canadensis</i> L.	116	41	59	perennial	herbaceous	native
Ranunculaceae	<i>Anemone hepatica</i> L.	95	40	60	perennial	herbaceous	native
Ranunculaceae	<i>Aquilegia canadensis</i> L.	119	251	238	perennial	herbaceous	native
Asteraceae	<i>Bidens vulgata</i> Greene	119	83	136	annual	herbaceous	native
Celastraceae	<i>Celastrus orbiculatus</i> Thunb.	103	151	220	perennial	herbaceous	introduced
Asteraceae	<i>Centaurea stoebe</i> Tausch	111	93	161	perennial	herbaceous	introduced
Asteraceae	<i>Cirsium arvense</i> (L.) Scop.	118	186	171	perennial	herbaceous	introduced
Asteraceae	<i>Cirsium discolor</i> (Muhl. ex Willd.) Spreng.	117	46	93	perennial	herbaceous	native
Geraniaceae	<i>Geranium maculatum</i> L.	119	489	513	perennial	herbaceous	native
Geraniaceae	<i>Geranium robertianum</i> L.	119	48	307	perennial	herbaceous	native
Xanthorrhoeaceae	<i>Hemerocallis fulva</i> (L.) L.	115	144	45	perennial	herbaceous	introduced
Malvaceae	<i>Hibiscus moscheutos</i> L.	119	105	100	perennial	herbaceous	native
Balsaminaceae	<i>Impatiens capensis</i> Meerb.	120	153	501	annual	herbaceous	native
Iridaceae	<i>Iris pseudacorus</i> L.	117	90	66	perennial	herbaceous	introduced
Iridaceae	<i>Iris versicolor</i> L.	119	344	185	perennial	herbaceous	native
Liliaceae	<i>Lilium canadense</i> L.	117	139	27	perennial	herbaceous	native
Caprifoliaceae	<i>Lonicera</i> × <i>bella</i> Zab.	107	37	62	perennial	woody	introduced
Caprifoliaceae	<i>Lonicera canadensis</i> Bartram ex Marshall	120	194	201	perennial	woody	native
Caprifoliaceae	<i>Lonicera japonica</i> Thunb.	116	329	115	perennial	woody	introduced
Rosaceae	<i>Malus pumila</i> Mill.	118	74	40	perennial	woody	introduced
Malvaceae	<i>Malva neglecta</i> Wallr.	116	25	140	perennial	herbaceous	introduced
Onagraceae	<i>Oenothera perennis</i> L.	120	194	214	perennial	herbaceous	native
Orobanchaceae	<i>Orobanche uniflora</i> L.	118	213	105	annual	herbaceous	native
Rosaceae	<i>Rosa gallica</i> L.	108	45	17	perennial	woody	introduced
Rosaceae	<i>Rubus odoratus</i> L.	120	176	318	perennial	woody	native
Sarraceniacae	<i>Sarracenia purpurea</i> L.	119	234	75	perennial	herbaceous	native
Iridaceae	<i>Sisyrinchium mucronatum</i> Michx.	117	86	157	perennial	herbaceous	native
Solanaceae	<i>Solanum rostratum</i> Dunal	115	21	85	annual	herbaceous	native
Melanthiaceae	<i>Trillium grandiflorum</i> (Michx.) Salisb.	119	129	40	perennial	herbaceous	native
Melanthiaceae	<i>Trillium undulatum</i> Willd.	120	402	156	perennial	herbaceous	native

and scored as fruiting if they had less than 50% flowers and buds and at least one fruit present. We used the *Thoreau's Field Notes* instance of *CrowdCurio* to crowdsourc phenological data from digitized herbarium specimens [46]. Citizen scientists hired through Amazon's Mechanical Turk service (MTurk; <https://www.mturk.com/>) counted the number of buds, flowers and fruits observed for a set of 10 specimen images. Participants first watched a short (1 min) instructional video on how to score phenological traits using *CrowdCurio* and then were provided with three tutorial images of each reproductive structure for every species. The 2364 anonymous participants were compensated at a rate of \$0.12 per image.

To provide an estimate of measurement error, each 10-image set scored by a single crowdsourcer included nine unique images and a single duplicate image randomly selected from the other nine [55]. We estimated the reliability score for each participant based on the data for each 10-image set by dividing the absolute

difference in organ counts for each phenophase by the total count of that specimen across the two duplicate specimens and subtracting this value from 1 ($1 - (|\text{count1} - \text{count2}| / (\text{count1} + \text{count2}))$) [55]. Reliability scores range from zero (unreliable/inconsistent) to one (reliable/consistent). Participants who reported no organs on one sheet and a non-zero number of the same organ on the duplicate sheet were assigned a reliability score of zero for that organ (i.e. the lowest reliability score). We conservatively selected the lowest reliability score among the three calculated for each organ per participant and assigned it to each participant as their final score. That is, if a participant got a reliability score equal to zero on one organ, they would be assigned a reliability score of zero for all organs. Specimen observations scored by crowdsourcers with a reliability score of zero were excluded from the analysis. We also spot checked for suspicious outliers manually and removed such data.

(c) Historical climate data

We used estimates of historic (1895–2016) average monthly temperature and precipitation data at 4 km resolution from PRISM (product AN81 m; <http://prism.oregonstate.edu/>), which provide high-resolution time-series estimates of climatic elements for the contiguous United States. As accurate locality data are not available for the majority of historic specimen records [59], we used county as our geographical unit of analysis. The vast majority (79%) of specimens used in this study were collected before 1980, and while 72% of the specimens used in this study had associated coordinate data, at least 91% of those coordinates had been georeferenced *post hoc* (e.g. assigned county or township centroid coordinates), and thus may not represent precise sampling locales. For each county and year, we estimated the mean monthly temperature, precipitation and elevation, and assigned these values to each specimen [59]. Although counties can vary in size and climate, counties in states along the Atlantic coast of the United States are generally small in size and geographically homogeneous. We estimated within-county climatic heterogeneity as the standard deviation of estimated monthly climatic values across each county and year and included it in our initial analyses, but coefficients had Bayesian credibility intervals that were not credibly different from zero, so we dropped these terms from our final models.

(d) Statistical analyses

Phenological sensitivity to spring temperature—defined as the mean of March, April and May temperatures [46]—was defined as the slope of the linear relationship between the day of year (DOY) of a phenophase and the spring temperature of the corresponding location and year (shifts in days per degree Celsius change: days/°C) [44,46]. These months have been used to define spring across the east coast of the United States [46,60]. To calculate phenological sensitivity, we binned our specimen data into both broad climatic zones [61,62] and finer-scale National Ecological Observatory Network (NEON) domains. Our data comprised two climatic zones (cold/very cold; mixed-humid/hot-humid—hereafter referred to as cold and mixed-warm) and five NEON ecoclimatic domains (NE, northeast; MA, mid-Atlantic; AP, Appalachians & Cumberland Plateau; OZ, Ozarks Complex; SE, southeast; electronic supplementary material, table S2). We also estimated phenological sensitivity to elevation as the slope of the linear relationship between the DOY of a phenophase and metres above sea level (m a.s.l.).

We estimated the mean timing of flowering and fruiting phenophases, and environmental influences on them, using Bayesian hierarchical linear regression models [63]. In our models, species, region, county and observer were considered random effects, while spring temperature and county elevation were covariates. The hierarchical nature of the model, in which the phenological responses of individual species were assumed to be drawn from statistical distributions instead of fixed estimates [64], allowed us to better estimate their climatic sensitivities. These models also more accurately quantified uncertainty in our estimates and partitioned the variance in phenological timing and phenological sensitivity within and between species and regions.

We fitted two models for each phenological phase. ‘Model 1’ estimated species-specific phenological sensitivities and partitioned their variances. ‘Model 2’ provided a more powerful comparison between phenological sensitivities found in the warmer, lower latitudes of our study area and those in the cooler, higher-latitude regions (figure 1).

In Model 1, the dependent variable was the DOY for which a given phenological phase (flowering or fruiting) was recorded for the i th specimen. $DOY_{[i]}$ was assumed to be normally distributed, with mean $\mu_{[i]}$ and species-specific variance $\sigma_{[s]}$.

$$DOY_{[i]} \sim N(\mu_{[i]}, \sigma_{[s]}). \quad (2.1)$$

The linear predictor $\mu_{[i]}$ was estimated as a function of covariates, including mean spring (March–May) air temperatures ($SpringT_{[i]}$) and the average elevation of the county in which the specimen was recorded ($Elev_{[c]}$). Additional intercept terms (α_1 – α_5) were added for each species (s), region (r), species \times region combination (sr), county (c) and observer (o). The full expression for estimating $\mu_{[i]}$ was

$$\begin{aligned} \mu_{[i]} = & \alpha_1_{[s]} + \alpha_2_{[r]} + \alpha_3_{[sr]} + \alpha_4_{[c]} + \alpha_5_{[o]} + \beta_1_{[s]} * SpringT_{[i]} \\ & + \beta_2_{[r]} * SpringT_{[i]} + \beta_3_{[sr]} * SpringT_{[i]} + \beta_4_{[s]} * Elev_{[c]}. \end{aligned} \quad (2.2)$$

Species-specific slope and intercept terms ($\alpha_1_{[s]}$, $\beta_1_{[s]}$ and $\beta_4_{[s]}$ in equation (2.2)) were drawn from normal distributions, with species assemblage means μ_{α_1} , μ_{β_1} and μ_{β_4} , and hypervariances σ_{α_1} , σ_{β_1} and σ_{β_4} .

$$\alpha_1_{[s]} \sim N(\mu_{\alpha_1}, \sigma_{\alpha_1}), \quad (2.3)$$

$$\beta_1_{[s]} \sim N(\mu_{\beta_1}, \sigma_{\beta_1}) \quad (2.4)$$

$$\text{and } \beta_4_{[s]} \sim N(\mu_{\beta_4}, \sigma_{\beta_4}). \quad (2.5)$$

Region and species \times region slopes ($\beta_2_{[r]}$, $\beta_3_{[sr]}$), and region, species \times region, county and observer intercepts ($\alpha_2_{[r]}$, $\alpha_3_{[sr]}$, $\alpha_4_{[c]}$, $\alpha_5_{[o]}$) in equation (2.2) were drawn from zero-centred normal distributions with hypervariances σ_{β_2} , σ_{β_3} , σ_{α_2} , σ_{α_3} , σ_{α_4} and σ_{α_5} , respectively. The species-specific sampling variation ($\sigma_{[s]}$) terms in equation (2.1) were estimated independently to account for differences in the duration of flowering and fruiting phases between species.

The three different groups of slopes estimated for spring temperature decomposed variation in phenological sensitivity into components representing between-species variability ($\beta_1_{[s]}$), between-region variability ($\beta_2_{[r]}$) and within-species variability across regions ($\beta_3_{[sr]}$). The accompanying hypervariances (σ_{β_1} , σ_{β_2} , σ_{β_3}) directly represented these different sources of variability; comparing their relative magnitudes quantified the contributions of each source of variation to overall variation in phenological sensitivity. This model structure also provided estimates of the contributions of species turnover to differences in phenological sensitivity between regions. We estimated these contributions by analysing the output of Model 1, computing phenological sensitivities for each observation for each iteration of our model. We then used the mean and standard deviation of these estimates for each region to create region-specific estimates of mean phenological sensitivities and their variability. We assessed the contribution of community turnover by comparing estimates that included all three climate sensitivity terms ($\beta_1_{[s]} + \beta_2_{[r]} + \beta_3_{[sr]}$) with estimates that included only the terms that represent species-level variability in climate sensitivity ($\beta_1_{[s]}$). This strategy allowed us to infer what the mean phenological sensitivities would be across regions in the hypothetical case that they differed only in species composition, and species responded identically to climate across their ranges.

Model 2 differed from Model 1 in treating the region term ($\beta_2_{[r]}$) as a two-level fixed effect representing the climatic region from which the specimen was drawn.

$$\begin{aligned} \mu_{[i]} = & \alpha_2_{[r]} + \alpha_3_{[sr]} + \alpha_4_{[c]} + \alpha_5_{[o]} + \beta_2_{[r]} * SpringT_{[i]} \\ & + \beta_3_{[sr]} * SpringT_{[i]} + \beta_4_{[s]} * Elev_{[c]}. \end{aligned} \quad (2.6)$$

Model 2 maximized statistical power to compare overall phenological sensitivities of species between warmer, more southerly parts of our study area and cooler, more northerly areas. Instead of treating region-specific slopes and intercepts as normally distributed random grouping factors, we represented species \times region, county and observer terms as zero-centred and normally distributed. This structure allowed more direct inference about overall differences in phenological sensitivity between cool and




Figure 1. Distribution of herbarium specimens across geographical and environmental space, colour-coded by NEON domain (*a,c*) or climate region (*b,d*). Panels (*a*) and (*c*) show specimens in flower, and panels (*b*) and (*d*) show specimens in fruit. Specimens are referenced to county centroids, and jitter has been added to coordinates to reduce over-plotting. Average spring (March–May) air temperatures are strongly negatively correlated with latitude (Pearson correlation $r = -0.92$, panel *c*), but elevation and latitude are largely independent (panel *d*).

mixed-warm climate regions (differences between estimates of $\beta_{2[r]}$ for the two regions). Unlike Model 1, however, Model 2 lacks species-specific parameters ($\alpha_{1[s]}$ and $\beta_{1[s]}$) and did not provide species-specific estimates of phenological sensitivity or permit comparison of variability in phenological sensitivity within and between species.

We estimated all parameters of the two models using Hamiltonian Monte Carlo (HMC) [65] implemented in Stan (v. 2.17.3) [66] called from the *rstan* interface [67] in R [68]. HMC is a form of Markov chain Monte Carlo (MCMC) that efficiently estimates hierarchical Bayesian regression models for larger datasets like ours [69]. We used relatively uninformative prior distributions: zero-centred normal priors for slopes and intercepts, and truncated normal distributions for variances and hypervariances. To account for sampling behaviour and simplify prior choices, we scaled and centred the response variable $\text{DOY}_{[t]}$ and all continuous predictors by subtracting the mean and dividing by the standard deviation of each variable. For each model run for each phenophase, we estimated parameters using four MCMC chains of 4000 iterations each and discarded the first 2000 iterations of each chain (as burn-in). We assessed convergence of each model both visually and with the Gelman–Rubin statistic ($\hat{r} < 1.1$ for all parameters). We also assessed good model fit using visual posterior predictive checks implemented in the *bayesplot* R package [70]. All parameter estimates were based on at least 1000 effective posterior samples. Estimates reported in the results were back-

transformed to the original data scale to facilitate illustration and interpretation.

Code and data for reproducing these analyses are archived by Harvard Forest [71].

3. Results

Our focal species spanned wide geographical and climatic space (figure 1). They demonstrated diverse patterns of phenology and significant variation in responses to climate across species and geographies. Using Model 1, estimated mean (non-leap-year) flower timing at 7.4°C and 216 m a.s.l. (mean collection conditions for the specimens) varied from 10 May (Day 130, *Anemone hepatica*) to 10 September (Day 253, *Bidens vulgata*) for flowering and 22 May (Day 142, *A. hepatica*) to 14 September (Day 257, *B. vulgata*) for fruiting (figure 2). The average lag time between flowering and fruiting across all species was approximately 20 days. Most species flowered and fruited earlier with warmer spring temperatures (assemblage mean -2.56 days/ $^{\circ}\text{C}$, 95% CI -3.64 to -1.48 , figure 2), and these responses were credibly different from zero (posterior probability > 0.95) for 21 out of 30 species for flowering and 15 out of 30 species for fruiting (electronic supplementary material, tables S3 and S4).




Figure 2. Mean flowering and fruiting time compared with estimated phenological sensitivities to spring (March–May) temperatures (*a,b*), and collection elevation (*c,d*) of 30 species estimated from herbarium specimens using a Bayesian hierarchical model (Model 1). Coloured and enumerated dots indicate species-specific estimates, and white circles at panel margins represent estimated community means for each quantity. Thick and thin bars represent 50 and 80% posterior credible intervals, respectively. Thick black lines represent credible linear relationships between quantities on *x*- and *y*-axes (posterior slope estimate different from zero with greater than 90% probability). Dotted lines represent non-credible relationships (posterior slope estimate not different from zero with greater than 90% probability).

For both flowering and fruiting, species with earlier reproductive phenologies were substantially more sensitive to spring temperature than species that flowered and fruited later in the season. This sensitivity manifested in a strong positive correlation between mean flowering and fruiting date and spring temperature sensitivity, with a slope of 0.018 days/°C per day for flowering and 0.023 days/°C per day for fruiting (figure 2*a,b*). These slopes were different from zero with greater than 99% posterior probability. We also found that, all other conditions being equal, flowering and fruiting came earlier at higher elevations (community means -1.71 to -0.11 days earlier per 100 m greater elevation for flowering, and -2.07 to 0.14 days earlier for fruiting, figure 2*c,d*). These effects were credibly different from zero for 7 of 30 species for flowering and 8 of 30 species for fruiting, but elevation influenced early-flowering and late-flowering species approximately equally.

Species in the warm and mixed-temperate climatic regions showed greater mean sensitivities to spring temperature and also greater variability (standard deviation) in climate sensitivity between species than in the cool-temperate northeast and Appalachians (figure 3; electronic supplementary material, figures S1 and S2). Using Model 2, we estimated that mean sensitivities in the mixed-warm region (figure 1*b*) were -2.96 days/°C (95% CI 3.69 to -2.25) for flowering and -3.37 days/°C (95% CI -4.12 to -2.60) for fruiting, but were substantially closer to zero in the cool-temperate region (-2.51 days/°C, 95% CI -2.86 to -2.19 for flowering, and -2.09 days/°C, 95% CI -2.63 to -1.57 for fruiting, figure 3*a*). The mixed-warm climatic region also had greater assemblage variability in phenological sensitivity

(figure 3*b*). All differences between cold and mixed-warm climatic regions had a posterior probability greater than 0.95 except for mean differences in flowering, where differences had a posterior probability of 0.87. These qualitative patterns were robust to an alternative spatial binning strategy that used only latitude, and not climate, to differentiate more northerly and southerly regions (electronic supplementary material, figure S3).

Overall differences between cool, northern and warm, southerly parts of the study area were accompanied by large amounts of regional variation not explained by climate or latitude (figure 4). For example, using Model 1 we estimated that plants in the Ozark Complex and mid-Atlantic NEON domains were substantially more phenologically sensitive than those in the northeast and Appalachians (figure 4*a*), while the northeast and Ozark Complex had plant assemblages with greater variability in phenological sensitivity (figure 4*c*). Differences in phenological sensitivity between regions were not adequately explained by differences in species composition, as per-sample weighted means computed using only species effects ($\beta_{1[s]}$ in equation (2.2)), did not show strong regional differences (figure 4*b,d*).

Our hierarchical approach allowed us to compare within-species, between-species and between-region sources of variability for both mean flowering time and sensitivity to spring temperature (figure 5). This analysis shows that between-species variation dominated variability in mean flowering time (figure 5*a*), but there was a similar amount of variation in phenological sensitivity within species between regions to that seen between species for both flowering and fruiting (figure 5*b*).




Figure 3. Estimated community phenological sensitivities to spring (March–May) air temperature in cold-temperate versus mixed-warm temperate climate zones (depicted in figure 1) using a Bayesian hierarchical model (Model 2). Panel (a) depicts species assemblage means, and panel (b) depicts assemblage standard deviations. Black and grey represent flowering and fruiting stages, and thick and thin bars represent 50 and 80% posterior credible intervals, respectively. Comparisons with a posterior probability greater than or equal to 0.8 and less than 0.95 are depicted with a ‘#’, comparisons with probabilities greater than or equal to 0.95 and less than 0.99 are depicted with ‘*’, and comparisons with probabilities greater than or equal to 0.99 are depicted with ‘***’.

4. Discussion

Our analyses revealed that (i) plant species from the eastern United States exhibit advanced timing of flowering and fruiting in response to warmer spring temperatures, (ii) the magnitude of these responses varies significantly between and within species across their latitudinal ranges and (iii) that phenological sensitivity to temperature tends to be higher in the warmer, more stable climates of lower-latitude regions.

(a) Differential responses to spring warming across species

Consistent with previous field observations of community phenology, we found that reproductive phenology of flowering plants accelerated with warming spring temperatures (e.g. [46,72,73]; but see [63]). The average number of days of phenological advancement per degree increase in temperature (-2.56 days/°C) that we observed also fell within previous estimates [46,74,75]. All else being equal, flowering and fruiting tended to occur earlier at higher elevations. Higher elevations tend to be relatively colder and have shorter

growing seasons, which exert pressure for species to initiate growth as soon as conditions become favourable [44,76–78].

Despite these general trends, we observed significant variation among species in their responses to warming. In general, early-flowering and early-fruiting species were more sensitive to spring temperatures than late-flowering/fruiting species (figure 2), a pattern also observed at smaller scales [75,79]. Warming-induced leaf budburst advancement has been suggested to be less prominent in late-flushing species compared with early-flushing ones owing to their greater chilling requirements [80]. Similar mechanisms may affect flowering and fruiting, where advances in the flowering date of late-flowering species caused by spring warming would be smaller than those of early-flowering species, which would manifest as weaker phenological responses to temperature in late-flowering species. Further, as flowering and fruiting events later in the year are more separated from spring climatic conditions, there is an extended window of time in which other factors could affect or modify reproductive timing. For instance, late-flowering species may be more sensitive to photoperiod or precipitation.

A large amount of variability in phenological sensitivity across species suggests that phenological responses to climatic change will be heterogeneous within communities. This could cause temporal reorganization of the structure and composition of plant communities, potentially altering direct and indirect interactions among plant species and between plant and animal species, and ecosystem services [24,34,81–83].

(b) Phenological sensitivity to spring temperature tends to decrease with latitude

The consequences of phenological shifts can be further complicated by intraspecific variation in phenological sensitivity to environmental cues [33,38]. For instance, using 20 years of observational data, Prevéy *et al.* [44] found that the phenological sensitivity to temperature of tundra plants at colder, higher latitudes was greater than at warmer, lower latitudes. However, contrary to such studies, we found that plants in warmer, lower-latitude regions tended to be more phenologically sensitive to temperature, especially for fruiting (figure 3). We hypothesize that this is due to the lower and less predictable winter and spring climates of the northeastern United States. In such environments, dynamic phenological tracking of spring temperatures (i.e. high phenological sensitivity to temperature) presents high risks to reproductive success, because warm periods may often be followed by periods of dramatic chilling [40]. At lower latitudes, the advent and progression of spring is less variable and average temperatures are higher; thus phenological tracking of temperature is less risky (electronic supplementary material, figures S1 and S2). Indeed, species exhibited a larger amount of variability in their responses to temperature in the warmer, lower-latitude parts of their ranges.

Climate and phenology might play different roles in filtering species assemblages in regions with longer growing seasons than in regions where the growing season is short and reproductive phenologies are strongly constrained by shorter freeze-free periods [35]. Indeed, studies synthesizing plot-level observational data have suggested phenological sensitivity of plant communities to warming may be positively correlated with mean annual temperature, but




Figure 4. Differences in sensitivity to spring temperatures between NEON domains are driven by regional variation due to local adaptation or plasticity, not community turnover between regions. This applies to both differences in species assemblage mean phenological sensitivities (*a,b*) and assemblage standard deviations (*c,d*). Panels (*a*) and (*b*) represent best estimates of regional variations in phenological sensitivity incorporating species identity, NEON Domain and NEON Domain \times species identity as random effects ($\beta_{1[st]}$, $\beta_{2[st]}$, $\beta_{3[st]}$) and panels (*c*) and (*d*) represent estimates incorporating only species-level effects ($\beta_{1[st]}$). This means that in panels (*c*) and (*d*), phenological sensitivity is assumed to be constant within a species across domains. Estimates for flowering are represented in black and estimates for fruiting are represented in grey. Thick and thin bars represent 50 and 80% posterior credible intervals, respectively, from a hierarchical Bayesian model (Model 1).

negatively correlated with seasonal temperature range (i.e. variability) in Europe [84] and China [85]. Alternatively, it is possible that plants in warmer climates exist closer to their response thresholds in terms of phenology, and thus react more dramatically to small changes in temperature. However, Körner & Basler [86] noted that cherry cultivars from regions with less variable spring temperatures flowered earlier in common gardens, suggesting phenological sensitivity does vary with climate. Plants in regions with high spring temperature variability also tend to be less phenologically sensitive in terms of leaf out and bud break to temperature than those in less variable climates [40]. Our results demonstrating that phenological sensitivity to temperature is higher in areas with low standard deviation of intra-annual temperature and inter-annual variation in spring temperature and high mean annual temperature support these findings.

(c) Consequences of variation in phenological responses across species ranges

Our results imply that with equal warming, individuals in lower-latitude populations will advance their reproductive phenology more dramatically than those at higher latitudes. This observed variation in phenological response may reflect adaptation to local climatic conditions, especially in annual species. We found a large amount of regional variation in

phenological sensitivity that was not clearly linked to climate or latitude. These regional differences were not explained by species turnover, but rather suggest the presence of inter-population variation driven by local adaptation or phenological plasticity (figure 4). Further, within-species variation in phenological sensitivity between regions was of similar magnitude to differences in sensitivity between species (figure 5). Other studies examining leaf out and senescence in trees also have shown that individuals from geographically and climatically separated populations differ in their phenology even when grown in common gardens [40,87,88].

Because the eastern United States is experiencing geographically variable climatic change, the heterogeneity in phenological responses to warming that we observed within and among species may have important consequences for plant communities in the near future. Colder, climatically variable high-latitude regions are experiencing disproportionate warming and climatic homogenization (i.e. reduced standard variation of intra-annual temperature), while warmer, climatically less variable more southerly regions are experiencing increases in intra-annual temperature variability (electronic supplementary material, figure S1). These climatic changes could alter patterns of overlap in reproductive timing among species in a community and across their individual ranges. Changes in phenological overlap across ranges could have direct consequences for adaptive evolution and species resilience to current and impending climatic changes, as




Figure 5. Variance components from Model 1, allowing comparisons of variability between NEON domains, between species and within species. Variance in the intercept (a) represents variability in flowering (black) or fruiting times (grey) at 7.4°C spring temperature and 216 m elevation above sea level, median conditions for the specimens. Variance in phenological sensitivity to spring air temperature (days/°C) represents variability in the slope of the linear relationship between spring temperature and flowering or fruiting times.

gene flow may increase among previously (temporally) isolated populations of some species and decrease among others [41–43,89–91]. Decreased phenological overlap could genetically isolate fringe populations, potentially leading to local extirpation and species range contractions. Moreover, phenologically sensitive plants at lower latitudes may especially be at risk due to increasingly variable temperatures and increased probabilities of phenological mismatch with mutualists [92]. However, we cannot fully rule out the possibility that individuals of long-lived perennial species may also be able to acclimate their phenological sensitivity to changing climatic conditions over longer periods of time.

Some of the variation we observe in phenological sensitivity to temperature across and within species may be due to differences in microclimate. However, the lack of accurate location data for most historic specimens limited our ability to infer fine-scale climate, necessitating coarser, county-level analyses. Also, we cannot ignore the possibility that the phenological trends we observed are unique to the species that we studied and/or reflect biases in herbarium collections [93]. To minimize effects of spatial bias and uncertainty, we studied common, well-collected species and accounted for climatic heterogeneity present in each sampling locale in our models. However, these taxa are not necessarily representative of species assemblages across regions, and our analyses do not account explicitly for spatial and temporal sampling

biases; some regional differences could be due to differing patterns of collection across space and time. Including county-level random effects as we have done here minimizes the impacts of these biases but does not eliminate them altogether. It is possible that different patterns of phenological sensitivity may be observed across species ranges depending on how climatic and/or geographical regions are delimited, though testing an alternative threshold yielded similar results, suggesting that the patterns we observe are robust to spatial binning choices. Additionally, we addressed crowdsourcing bias by including crowdsourcer random effects and removing observations for crowdsourcers with low reliability scores even though phenological data collected by citizen scientists do not differ significantly in quality from those collected by experts [46,55]. Lastly, although spring temperature is a critical driver of flowering phenology in temperate climates, we cannot fully exclude the possibility that other variables correlated with latitude or mean spring temperature may determine observed variance in phenological sensitivity [46,73,75,94–97]. For example, spring temperature tends to be highly correlated with mean annual and mean monthly temperatures in eastern North America (electronic supplementary material, figure S4). Photoperiod or snow melt may further influence and alter species phenological responses [98–103]. Future research into how these environmental cues interact to trigger phenological events is necessary and will greatly improve our understanding of plant phenology.

5. Conclusion

Building on previous phenological research by scoring multiple phenological traits across over 8000 herbarium specimens spanning 120 years, we have demonstrated that phenological sensitivity can vary greatly across species' ranges. This variance may be attributed to adaptation or acclimation to local climates. The large amount of within-species variation in phenological sensitivity that we observed underlines the complex and contingent nature of phenological sensitivities. Phenological responses of individual species to climate are not stable phenotypic traits, but instead emerge from a multitude of potentially reciprocal interactions between organisms and their environment. Populations in different regions could have differences in frequencies of genes that control how climate affects the timing of development or differences in microhabitat distributions between regions that alter how populations experience local climate. The regions themselves may have differences in unmeasured environmental factors that interact with responses to temperature or differences in species interactions that may alter phenological signalling. The circumstances and extent to which these or other factors explain regional variation in species responses to climate is currently unknown. To untangle the roles of ecological and evolutionary processes governing the heterogeneous phenological responses of plant species to warming, researchers will have to take advantage of new techniques and datasets. In addition to continued field observations and laboratory analysis of mechanisms responsible for flowering and fruiting, herbarium specimens can provide a comprehensive, nuanced picture of phenological responses to ongoing climatic change across many species. Our study further demonstrates that we can now harness the treasure trove of

information in herbaria across the world to examine hundreds, if not thousands of species across myriad plant lineages, habitats and regions. Such efforts will be critical to enhance our ability to forecast future changes in plant assemblages across space and time in an era of accelerating climate change [104,105].

Ethics. The use and collection of data by citizen scientists were approved by an ethics review committee at the University of Waterloo (ORE no. 21647).

Data accessibility. Data and R code are available from the Harvard Forest Data Archive (<http://harvardforest.fas.harvard.edu/dataarchive>, dataset HF309).

Authors' contributions. C.C.D. conceived the study; D.S.P., C.C.D., A.C.W. and E.L. designed the study; D.S.P. and A.C.W. collected data; D.S.P.,

A.M.E. and I.B. analysed the data; D.S.P. drafted the first version of the manuscript; C.C.D. and D.S.P. made first substantial revisions to this draft, and all authors contributed to subsequent revisions.

Competing interests. We declare we have no competing interests.

Funding. This study was funded as part of the New England Vascular Plant Project to C.C.D. (NSF-DBI: EF1208835), NSF-DEB 1754584 to C.C.D. and A.M.E., and an NSERC Discovery Grant to E.L. (RGPIN-2015-04543). A.M.E.'s participation in this project was supported by Harvard Forest. D.S.P.'s contribution was supported by the Harvard University Herbaria and NSF-DEB 1754584. I.B.'s contribution was supported by the National Science Foundation's Postdoctoral Research Fellowship in Biology (NSF-DBI-1711936).

Acknowledgements. The authors thank X. Feng, S. Worthington, and members of the Davis lab for their invaluable insights and comments on the project and manuscript.

References

- Field CB *et al.* 2014 Climate change 2014: impacts, adaptation, and vulnerability. Part A: global and sectoral aspects. In *Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Geneva, Switzerland: IPCC.
- Root TL, Price JT, Hall KR, Schneider SH, Rosenzweig C, Pounds JA. 2003 Fingerprints of global warming on wild animals and plants. *Nature* **421**, 57–60. (doi:10.1038/nature01333)
- Parnesan C, Yohe G. 2003 A globally coherent fingerprint of climate change impacts across natural systems. *Nature* **421**, 37–42. (doi:10.1038/nature01286)
- Miller-Rushing AJ, Primack RB, Primack D, Mukunda S. 2006 Photographs and herbarium specimens as tools to document phenological changes in response to global warming. *Am. J. Bot.* **93**, 1667–1674. (doi:10.3732/ajb.93.11.1667)
- Menzel A, Sparks TH, Estrella N, Roy DB. 2006 Altered geographic and temporal variability in phenology in response to climate change. *Glob. Ecol. Biogeogr.* **15**, 498–504. (doi:10.1111/j.1466-822X.2006.00247.x)
- Chambers LE *et al.* 2013 Phenological changes in the southern hemisphere. *PLoS ONE* **8**, e75514. (doi:10.1371/journal.pone.0075514)
- Ovaskainen O, Skorokhodova S, Yakovleva M, Sukhov A, Kutenkov A, Kutenkova N, Shcherbakov A, Meyke E, del Delgado M. 2013 Community-level phenological response to climate change. *Proc. Natl Acad. Sci. USA* **110**, 13 434–13 439. (doi:10.1073/pnas.1305533110)
- Willis CG, Ruhfel B, Primack RB, Miller-Rushing AJ, Davis CC. 2008 Phylogenetic patterns of species loss in Thoreau's woods are driven by climate change. *Proc. Natl Acad. Sci. USA* **105**, 17 029–17 033. (doi:10.1073/pnas.0806446105)
- Inouye D. 2008 Effects of climate change on phenology, frost damage, and floral abundance of montane wildflowers. *Ecology* **89**, 353–362. (doi:10.1890/06-2128.1)
- Wheeler HC, Høye TT, Schmidt NM, Svenning JC, Forchhammer MC. 2015 Phenological mismatch with abiotic conditions—implications for flowering in Arctic plants. *Ecology* **96**, 775–787. (doi:10.1890/14-0338.1)
- White MA, Nemani RR. 2003 Canopy duration has little influence on annual carbon storage in the deciduous broad leaf forest. *Glob. Change Biol.* **9**, 967–972. (doi:10.1046/j.1365-2486.2003.00585.x)
- Richardson AD, Hollinger DY, Dail DB, Lee JT, Munger JW, O'keefe J. 2009 Influence of spring phenology on seasonal and annual carbon balance in two contrasting New England forests. *Tree Physiol.* **29**, 321–331. (doi:10.1093/treephys/tpn040)
- Barford CC *et al.* 2001 Factors controlling long- and short-term sequestration of atmospheric CO₂ in a mid-latitude forest. *Science* **294**, 1688–1691. (doi:10.1126/science.1062962)
- Fitzjarrald DR, Acevedo OC, Moore KE. 2001 Climatic consequences of leaf presence in the eastern United States. *J. Clim.* **14**, 598–614. (doi:10.1175/1520-0442(2001)014<0598:CCOLPI>2.0.CO;2)
- Inouye DW, Barr B, Armitage KB, Inouye BD. 2000 Climate change is affecting altitudinal migrants and hibernating species. *Proc. Natl Acad. Sci. USA* **97**, 1630–1633. (doi:10.1073/pnas.97.4.1630)
- Visser ME, Both C. 2005 Shifts in phenology due to global climate change: the need for a yardstick. *Proc. R. Soc. B* **272**, 2561–2569. (doi:10.1098/rspb.2005.3356)
- Doiron M, Gauthier G, Lévesque E. 2015 Trophic mismatch and its effects on the growth of young in an Arctic herbivore. *Glob. Change Biol.* **21**, 4364–4376. (doi:10.1111/gcb.13057)
- Post E, Forchhammer MC. 2008 Climate change reduces reproductive success of an Arctic herbivore through trophic mismatch. *Phil. Trans. R. Soc. B* **363**, 2367–2373. (doi:10.1098/rstb.2007.2207)
- Post E, Pedersen C, Wilmers CC, Forchhammer MC. 2008 Warming, plant phenology and the spatial dimension of trophic mismatch for large herbivores. *Proc. R. Soc. B* **275**, 2005–2013. (doi:10.1098/rspb.2008.0463)
- Plard F, Gaillard J-M, Coulson T, Hewison AJM, Delorme D, Wornant C, Bonenfant C. 2014 Mismatch between birth date and vegetation phenology slows the demography of roe deer. *PLoS Biol.* **12**, e1001828. (doi:10.1371/journal.pbio.1001828)
- Kharouba HM, Ehrlén J, Gelman A, Bolmgren K, Allen JM, Travers SE, Wolkovich EM. 2018 Global shifts in the phenological synchrony of species interactions over recent decades. *Proc. Natl Acad. Sci. USA* **115**, 5211–5216. (doi:10.1073/pnas.1714511115)
- Navarro-Cano JA, Karlsson B, Posledovich D, Toftegaard T, Wiklund C, Ehrlén J, Gotthard K. 2015 Climate change, phenology, and butterfly host plant utilization. *Ambio* **44**, 78–88. (doi:10.1007/s12280-014-0602-z)
- Posledovich D, Toftegaard T, Wiklund C, Ehrlén J, Gotthard K. 2018 Phenological synchrony between a butterfly and its host plants: experimental test of effects of spring temperature. *J. Anim. Ecol.* **87**, 150–161. (doi:10.1111/1365-2656.12770)
- Burkle LA, Marlin JC, Knight TM. 2013 Plant-pollinator interactions over 120 years: loss of species, co-occurrence, and function. *Science* **339**, 1611–1615. (doi:10.1126/science.1232728)
- Kudo G, Ida TY. 2013 Early onset of spring increases the phenological mismatch between plants and pollinators. *Ecology* **94**, 2311–2320. (doi:10.1890/12-2003.1)
- Iler AM, Inouye DW, Høye TT, Miller-Rushing AJ, Burkle LA, Johnston EB. 2013 Maintenance of temporal synchrony between syrphid flies and floral resources despite differential phenological responses to climate. *Glob. Change Biol.* **19**, 2348–2359. (doi:10.1111/gcb.12246)
- Visser ME, van Noordwijk AJ, Tinbergen JM, Lessells CM. 1998 Warmer springs lead to mistimed reproduction in great tits (*Parus major*). *Proc. R. Soc. Lond. B* **265**, 1867–1870. (doi:10.1098/rspb.1998.0514)
- Mayor SJ *et al.* 2017 Increasing phenological asynchrony between spring green-up and arrival of migratory birds. *Sci. Rep.* **7**, 1902. (doi:10.1038/s41598-017-02045-z)

29. Wolkovich EM, Cleland EE. 2011 The phenology of plant invasions: a community ecology perspective. *Front. Ecol. Environ.* **9**, 287–294. (doi:10.1890/100033)
30. Craine JM, Wolkovich EM, Towne EG, Kembel SW. 2012 Flowering phenology as a functional trait in a tallgrass prairie. *New Phytol.* **193**, 673–682. (doi:10.1111/j.1469-8137.2011.03953.x)
31. Colautti RI, Ågren J, Anderson JT, Colautti RI, Jon A, Colautti RI. 2017 Phenological shifts of native and invasive species under climate change: insights from the *Boechera*–*Lythrum* model. *Phil. Trans. R. Soc. B* **372**, 20160032. (doi:10.1098/rstb.2016.0032)
32. Pau S, Wolkovich EM, Cook BI, Davies TJ, Kraft NJB, Bolmgren K, Betancourt JL, Cleland EE. 2011 Predicting phenology by integrating ecology, evolution and climate science. *Glob. Change Biol.* **17**, 3633–3643. (doi:10.1111/j.1365-2486.2011.02515.x)
33. Wolkovich EM, Cook BI, Davies TJ. 2014 Progress towards an interdisciplinary science of plant phenology: building predictions across space, time and species diversity. *New Phytol.* **201**, 1156–1162. (doi:10.1111/nph.12599)
34. CaraDonna PJ, Iler AM, Inouye DW. 2014 Shifts in flowering phenology reshape a subalpine plant community. *Proc. Natl Acad. Sci. USA* **111**, 4916–4921. (doi:10.1073/pnas.1323073111)
35. Diez JM, Ibáñez I, Miller-Rushing AJ, Mazer SJ, Crimmins TM, Crimmins MA, Bertelsen CD, Inouye DW. 2012 Forecasting phenology: from species variability to community patterns. *Ecol. Lett.* **15**, 545–553. (doi:10.1111/j.1461-0248.2012.01765.x)
36. Banta JA, Ehrenreich IM, Gerard S, Chou L, Wilczek A, Schmitt J, Kover PX, Purugganan MD. 2012 Climate envelope modelling reveals intraspecific relationships among flowering phenology, niche breadth and potential range size in *Arabidopsis thaliana*. *Ecol. Lett.* **15**, 769–777. (doi:10.1111/j.1461-0248.2012.01796.x)
37. Wilczek AM *et al.* 2009 Effects of genetic perturbation on seasonal life history plasticity. *Science* **323**, 930–934. (doi:10.1126/science.1165826)
38. Matthews ER, Mazer SJ. 2016 Historical changes in flowering phenology are governed by temperature \times precipitation interactions in a widespread perennial herb in western North America. *New Phytol.* **210**, 157–167. (doi:10.1111/nph.13751)
39. Barros VR *et al.* 2014 Climate change 2014: impacts, adaptation, and vulnerability. Part B: regional aspects.. In *Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Geneva, Switzerland: IPCC.
40. Zohner CM, Renner SS. 2014 Common garden comparison of the leaf-out phenology of woody species from different native climates, combined with herbarium records, forecasts long-term change. *Ecol. Lett.* **17**, 1016–1025. (doi:10.1111/ele.12308)
41. Lenormand T. 2002 Gene flow and the limits to natural selection. *Trends Ecol. Evol.* **17**, 183–189. (doi:10.1016/S0169-5347(02)02497-7)
42. Alleaume-Benharira M, Pen IR, Ronce O. 2006 Geographical patterns of adaptation within a species' range: interactions between drift and gene flow. *J. Evol. Biol.* **19**, 203–215. (doi:10.1111/j.1420-9101.2005.00976.x)
43. Sexton JP, Strauss SY, Rice KJ. 2011 Gene flow increases fitness at the warm edge of a species' range. *Proc. Natl Acad. Sci. USA* **108**, 11 704–11 709. (doi:10.1073/pnas.1100404108)
44. Prevéy J *et al.* 2017 Greater temperature sensitivity of plant phenology at colder sites: implications for convergence across northern latitudes. *Glob. Change Biol.* **23**, 2660–2671. (doi:10.1111/gcb.13619)
45. Riihimäki M, Savolainen O. 2004 Environmental and genetic effects on flowering differences between northern and southern populations of *Arabidopsis lyrata* (Brassicaceae). *Am. J. Bot.* **91**, 1036–1045. (doi:10.3732/ajb.91.7.1036)
46. Willis CG *et al.* 2017 CrowdCurio: an online crowdsourcing platform to facilitate climate change studies using herbarium specimens. *New Phytol.* **215**, 479–488. (doi:10.1111/nph.14535)
47. Willis CG *et al.* 2017 Old plants, new tricks: phenological research using herbarium specimens. *Trends Ecol. Evol.* **32**, 531–546. (doi:10.1016/j.tree.2017.03.015)
48. Davis CC, Willis CG, Connolly B, Kelly C, Ellison AM. 2015 Herbarium records are reliable sources of phenological change driven by climate and provide novel insights into species' phenological cueing mechanisms. *Am. J. Bot.* **102**, 1599–1609. (doi:10.3732/ajb.1500237)
49. Primack D, Imbres C, Primack RB, Miller-Rushing AJ, Del Tredici P. 2004 Herbarium specimens demonstrate earlier flowering times in response to warming in Boston. *Am. J. Bot.* **91**, 1260–1264. (doi:10.3732/ajb.91.8.1260)
50. Robbitt KM, Davy AJ, Hutchings MJ, Roberts DL. 2011 Validation of biological collections as a source of phenological data for use in climate change studies: a case study with the orchid *Ophrys sphegodes*. *J. Ecol.* **99**, 235–241. (doi:10.1111/j.1365-2745.2010.01727.x)
51. Vellend M, Brown CD, Kharouba HM, McCune JL, Myers-Smith IH. 2013 Historical ecology: using unconventional data sources to test for effects of global environmental change. *Am. J. Bot.* **100**, 1294–1305. (doi:10.3732/ajb.1200503)
52. Sweeney PW, Starly B, Morris PJ, Xu Y, Jones A, Radhakrishnan S, Grassa CJ, Davis CC. 2018 Large-scale digitization of herbarium specimens: development and usage of an automated, high-throughput conveyor system. *Taxon* **67**, 165–178. (doi:10.12705/671.9)
53. Yost JM *et al.* 2018 Digitization protocol for scoring reproductive phenology from herbarium specimens of seed plants. *Appl. Plant Sci.* **6**, e1022. (doi:10.1002/aps3.1022)
54. Pearse WD, Davis CC, Inouye DW, Primack RB, Davies TJ. 2017 A statistical estimator for determining the limits of contemporary and historic phenology. *Nat. Ecol. Evol.* **1**, 1876. (doi:10.1038/s41559-017-0350-0)
55. Williams AC, Goh J, Willis CG, Ellison AM, Brusuelas JH, Davis CC, Law E. 2017 Déjà vu: characterizing worker reliability using task consistency. In *Proc. 5th AAAI Conf. Human Computation and Crowdsourcing, HCOMP 2017, 23–26 October 2017, Québec City, Québec, Canada* (eds S Dow, AT Kalai), pp. 197–205. Palo Alto, CA: AAAI Press.
56. Schorn C, Weber E, Bernardos R, Hopkins C, Davis C. 2016 The New England Vascular Plants Project: 295 000 specimens and counting. *Rhodora* **118**, 324. (doi:10.3119/15-34)
57. SouthEast Regional Network of Expertise and Collections. 2017 *SERNEC Data Portal*. See <http://serneportal.org/portal/index.php> (accessed 28 June 2018).
58. Hart R, Salick J, Ranjitkar S, Xu J. 2014 Herbarium specimens show contrasting phenological responses to Himalayan climate. *Proc. Natl Acad. Sci. USA* **111**, 10 615–10 619. (doi:10.1073/pnas.1403376111)
59. Park DS, Davis CC. 2017 Implications and alternatives of assigning climate data to geographical centroids. *J. Biogeogr.* **44**, 2188–2198. (doi:10.1111/jbi.13029)
60. Peñá JE, Duncan R, Browning H. 1996 Seasonal abundance of *Phyllocnistis citrella* (Lepidoptera: Gracillariidae) and its parasitoids South Florida citrus. *Environ. Entomol.* **25**, 698–702. (doi:10.1093/ee/25.3.698)
61. Baechler MC, Gilbride TL, Cole PC, Hefty MG, Ruiz K. 2010 Guide to determining climate regions by county. *Build. Am. Best Pract. Ser.* **7**, 1–45. (doi:10.2172/969181)
62. Zhang Y, Bielory L, Georgopoulos PG. 2014 Climate change effect on *Betula* (birch) and *Quercus* (oak) pollen seasons in the United States. *Int. J. Biometeorol.* **58**, 909–919. (doi:10.1007/s00484-013-0674-7)
63. Hobbs NT, Hooten MB. 2015 *Bayesian models: a statistical primer for ecologists*. Princeton, NJ: Princeton University Press.
64. Gelman A, Hill J. 2006 *Data analysis using regression and multilevel/hierarchical models*. Cambridge, UK: Cambridge University Press.
65. Hoffman MD, Gelman A. 2014 The No-U-turn sampler: adaptively setting path lengths in Hamiltonian Monte Carlo. *J. Mach. Learn. Res.* **15**, 1593–1623.
66. Carpenter B *et al.* 2017 Stan: a probabilistic programming language. *J. Stat. Softw.* **76**, 1–32. (doi:10.18637/jss.v076.i01)
67. Stan Development Team. 2018 RStan: the R interface to Stan. R package version 2.18.1. See <http://mc-stan.org/>.
68. R Core Team. 2018 *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing. See <http://www.R-project.org>.
69. Betancourt M, Girolami M. 2015 Hamiltonian Monte Carlo for hierarchical models. *Curr. Trends Bayesian Methodol. Appl.* **79**, 30. (doi:10.1201/b18502-5)
70. Gabry J, Simpson D, Vehtari A, Betancourt M, Gelman A. 2017 Visualization in Bayesian workflow. See <http://arxiv.org/abs/1709.01449>.

71. Park DS, Breckheimer I, Williams AC, Law E, Ellison AM, Davis CC. 2018 Assessing plant phenological patterns in the eastern United States over the last 120 years. Harvard Forest Data Archive, HF309. (doi:10.6073/pasta/f6afa728bb5edfd79f458d7d5e23f559)
72. Roberts AMI, Tansey C, Smithers RJ, Phillimore AB. 2015 Predicting a change in the order of spring phenology in temperate forests. *Glob. Change Biol.* **21**, 2603–2611. (doi:10.1111/gcb.12896)
73. Yu H, Luedeling E, Xu J. 2010 Winter and spring warming result in delayed spring phenology on the Tibetan Plateau. *Proc. Natl Acad. Sci. USA* **107**, 22 151–22 156. (doi:10.1073/pnas.1012490107)
74. Wolfe DW, Schwartz MD, Lakso AN, Otsuki Y, Pool RM, Shaulis NJ. 2005 Climate change and shifts in spring phenology of three horticultural woody perennials in northeastern USA. *Int. J. Biometeorol.* **49**, 303–309. (doi:10.1007/s00484-004-0248-9)
75. Miller-Rushing AJ, Primack RB. 2008 Global warming and flowering times in Thoreau's Concord: a community perspective. *Ecology* **89**, 332–341. (doi:10.1890/07-0068.1)
76. Bliss LC. 1962 Adaptations of arctic and alpine plants to environmental conditions. *Arctic* **15**, 117–144. (doi:10.14430/arctic3564)
77. Billings WD, Mooney HA. 1968 The ecology of arctic and alpine plants. *Biol. Rev.* **43**, 481–529. (doi:10.1111/j.1469-185X.1968.tb00968.x)
78. Oberbauer SF *et al.* 2013 Phenological response of tundra plants to background climate variation tested using the International Tundra Experiment. *Phil. Trans. R. Soc. B* **368**, 20120481. (doi:10.1098/rstb.2012.0481)
79. Fitter AH, Fitter RSR. 2002 Rapid changes in flowering time in British plants. *Science* **296**, 1689–1691. (doi:10.1126/science.1071617)
80. Murray MB, Cannell MGR, Smith RI. 1989 Date of budburst of fifteen tree species in Britain following climatic warming. *J. Appl. Ecol.* **26**, 693–700. (doi:10.2307/2404093)
81. Theobald EJ, Breckheimer I, HilleRisLambers J. 2017 Climate drives phenological reassembly of a mountain wildflower meadow community. *Ecology* **98**, 2799–2812. (doi:10.1002/ecy.1996)
82. Miller-Rushing AJ, Inouye DW. 2009 Variation in the impact of climate change on flowering phenology and abundance: an examination of two pairs of closely related wildflower species. *Am. J. Bot.* **96**, 1821–1829. (doi:10.3732/ajb.0800411)
83. Memmott J, Craze PG, Waser NM, Price MV. 2007 Global warming and the disruption of plant–pollinator interactions. *Ecol. Lett.* **10**, 710–717. (doi:10.1111/j.1461-0248.2007.01061.x)
84. Lapeña A, Henry H, Vuille M, Mower J. 2014 Climatic factors controlling plant sensitivity to warming. *Clim. Change* **122**, 723–734. (doi:10.1007/s10584-013-1010-2)
85. Zhang H, Yuan W, Liu S, Dong W, Fu Y. 2015 Sensitivity of flowering phenology to changing temperature in China. *J. Geophys. Res. Biogeosci.* **120**, 1658–1665. (doi:10.1002/2015JG003112)
86. Körner C, Basler D. 2010 Phenology under global warming. *Science* **327**, 1461–1462. (doi:10.1126/science.1186473)
87. Zohner CM, Benito BM, Fridley JD, Svenning J, Renner SS. 2017 Spring predictability explains different leaf-out strategies in the woody floras of North America, Europe and East Asia. *Ecol. Lett.* **20**, 452–460. (doi:10.1111/ele.12746)
88. Zohner CM, Renner SS. 2017 Innately shorter vegetation periods in North American species explain native–non-native phenological asymmetries. *Nat. Ecol. Evol.* **1**, 1655. (doi:10.1038/s41559-017-0307-3)
89. Savolainen O, Pyhäjärvi T, Knürr T. 2007 Gene flow and local adaptation in trees. *Annu. Rev. Ecol. Evol. Syst.* **38**, 595–619. (doi:10.1146/annurev.ecolsys.38.091206.095646)
90. Nagy ES, Rice KJ. 1997 Local adaptation in two subspecies of an annual plant: implications for migration and gene flow. *Evolution* **51**, 1079–1089. (doi:10.1111/j.1558-5646.1997.tb03955.x)
91. Fox GA. 2003 Assortative mating and plant phenology: evolutionary and practical consequences. *Evol. Ecol. Res.* **5**, 1–18.
92. Thomson JD. 2010 Flowering phenology, fruiting success and progressive deterioration of pollination in an early-flowering geophyte. *Phil. Trans. R. Soc. B* **365**, 3187–3199. (doi:10.1098/rstb.2010.0115)
93. Daru BH *et al.* 2018 Widespread sampling biases in herbaria revealed from large-scale digitization. *New Phytol.* **217**, 939–955. (doi:10.1111/nph.14855)
94. Richardson AD, Bailey AS, Denny EG, Martin CW, O'Keefe J. 2006 Phenology of a northern hardwood forest canopy. *Glob. Change Biol.* **12**, 1174–1188. (doi:10.1111/j.1365-2486.2006.01164.x)
95. Ellwood ER, Temple SA, Primack RB, Bradley NL, Davis CC. 2013 Record-breaking early flowering in the eastern United States. *PLoS ONE* **8**, 1–9. (doi:10.1371/journal.pone.0053788)
96. Augspurger CK. 2013 Reconstructing patterns of temperature, phenology, and frost damage over 124 years: spring damage risk is increasing. *Ecology* **94**, 41–50. (doi:10.1890/12-0200.1)
97. Güsewell S, Furrer R, Gehrig R, Pietragalla B. 2017 Changes in temperature sensitivity of spring phenology with recent climate warming in Switzerland are related to shifts of the pre-season. *Glob. Change Biol.* **23**, 5189–5202. (doi:10.1111/gcb.13781)
98. Bjorkman AD, Elmendorf SC, Beamish AL, Vellend M, Henry GHR. 2015 Contrasting effects of warming and increased snowfall on Arctic tundra plant phenology over the past two decades. *Glob. Change Biol.* **21**, 4651–4661. (doi:10.1111/gcb.13051)
99. Way DA, Montgomery RA. 2015 Photoperiod constraints on tree phenology, performance and migration in a warming world. *Plant. Cell Environ.* **38**, 1725–1736. (doi:10.1111/pce.12431)
100. Nakagawa H, Yamagishi J, Miyamoto N, Motoyama M, Yano M, Nemoto K. 2005 Flowering response of rice to photoperiod and temperature: a QTL analysis using a phenological model. *Theor. Appl. Genet.* **110**, 778–786. (doi:10.1007/s00122-004-1905-4)
101. Dunne JA, Harte J, Taylor KJ. 2003 Subalpine meadow flowering phenology responses to climate change: integrating experimental and gradient methods. *Ecol. Monogr.* **73**, 69–86. (doi:10.1890/0012-9615(2003)073[0069:SMFPRT]2.0.CO;2)
102. Wolkovich EM, Davies TJ, Schaefer H, Cleland EE, Cook BI, Travers SE, Willis CG, Davis CC. 2013 Temperature-dependent shifts in phenology contribute to the success of exotic species with climate change. *Am. J. Bot.* **100**, 1407–1421. (doi:10.3732/ajb.1200478)
103. Park IW, Mazer SJ. In press. Overlooked climate parameters best predict flowering onset: assessing phenological models using the elastic net. *Glob. Change Biol.* (doi:10.1111/gcb.14447)
104. Chuine I, Beaubien EG. 2001 Phenology is a major determinant of tree species range. *Ecol. Lett.* **4**, 500–510. (doi:10.1046/j.1461-0248.2001.00261.x)
105. Chuine I. 2010 Why does phenology drive species distribution? *Phil. Trans. R. Soc. B* **365**, 3149–3160. (doi:10.1098/rstb.2010.0142)