

Root Systems of Individual Plants Worldwide 2022

Shersingh Joseph Tumber-Dávila, H. Jochen Schenk, Enzai Du, and Robert B. Jackson

References for RSIP Data

- Abbott ML, Fraley L Jr, Reynolds TD (1991) Root profiles of selected cold desert shrubs and grasses in disturbed and undisturbed soils. *Environmental and Experimental Botany* 31: 165-178
- Adams ME (1967) A study of the ecology of *Acacia mellifera*. *A. seyal* and *Balanites aegyptiaca* in relation to land-clearing. *Journal of Applied Ecology* 4: 221-237
- Ahmet Birand H (1938) Untersuchungen zur Wasserökologie der Steppenpflanzen bei Ankara. *Jahrbücher für wissenschaftliche Botanik* 87: 93-172
- Albertson FW (1937) Ecology of mixed prairie in west central Kansas. *Ecological Monographs* 7: 481-547
- Alexandre DY, Ouédraogo SJ (1992) Variations in Root Morphology of *Faidherbia albida* in Relation to Soil and Agronomic Effects (World Agroforestry Centre).
- Anderson VL (1927) Studies of the vegetation of the English Chalk: V. The water economy of the chalk flora. *Journal of Ecology* 15: 72-129
- Antos JA, Halpern CB (1997) Root system differences among species: implications for early successional changes in forests of western Oregon. *American Midland Naturalist* 138: 97-108
- Ashton DH (1975) The root and shoot development of *Eucalyptus regnans* F. Muell. *Australian Journal of Botany* 23: 867-887
- Baitulin IO (1979) Kornevaja sistema rastenij aridnoj zony Kazakhstana. (Root systems of plants of the arid zone of Kazakhstan. In Russian). Nauka. Alma-Ata
- Baitulin IO (1996) Root research in natural plant communities of Kazakhstan. *Acta Phytogeographica Suecica* 81: 7-10
- Baitulin IO. ed (1993) Fitoékologicheskie issledovaniya v juzhnoj Gobi. (Phytoecological investigations in southern Gobi. In Russian with English summary). Gylim. Alma-Ata
- Bang-xing W (1991) Studies on the vertical structure of seasonal rain-forest in Xishuangbanna of Yunnan. *Acta Botanica Sinica* 33: 232-239
- Barbour MG, Major J. eds (1988) Terrestrial vegetation of California. (New expanded edition 1988). California Native Plant Society. Davis
- Batanouny KH, Abdel Wahab AM (1973) Eco-physiological studies on desert plants VIII. Root penetration of *Leptadenia pyrotechnica* (Forsk.) Decne. in relation to its water balance. *Oecologia* 11: 151-161
- Batanouny KH, Batanouny MH (1969) Formation of phytogenic hillocks. II. Rooting habit of plants forming phytogenic hillocks. *Acta Botanica Academiae Scientiarum Hungaricae* 15: 1-18
- Becker P, Castillo A (1990) Root architecture of shrubs and saplings in the understory of a tropical moist forest in lowland Panama. *Biotropica* 22: 242-249

Bendali F, Floret C, Le Floc'h E, Ponantier R (1990) The dynamics of vegetation and sand mobility in arid regions of Tunisia. *Journal of Arid Environments* 18: 21-32

Berndt HW, Gibbons RD (1958) Root distribution of some native trees and understory plants growing on three sites within ponderosa pine watersheds in Colorado. US Dep. Agric. For. Serv. Sta. Pap. RM-37.

Bertiller MB, Beeskow AM, Coronato F (1991) Seasonal environmental variation and plant phenology in arid Patagonia (Argentina). *Journal of Arid Environments* 21: 1-11

Bhimaya CP, Kaul RN (1965) Root system of four desert tree species. *Annals of Arid Zone* 4: 185-194

Biswell HH (1935) Effects of environment upon the root habits of certain deciduous forest trees. *Botanical Gazette* 96: 676-708

Blagoveshchenskiy EN (1968) The dry savanna of northwest India. *Soviet Geography* 9: 519-537

Bonham CD Mack SE (1990) Root distributions of *Eurotia lanata* in association with two species of *Agropyron* on disturbed soils. *Botanical Gazette* 151: 522-527

Bowns JE, Jr, Box TW (1964) The influence of grazing on the roots and rhizomes of seacoast bluestem. *Journal of Range Management* 17: 36-39

Branson FA, Miller RF, McQueen IS (1976) Moisture relationships in twelve northern desert shrub communities near Grand Junction. Colorado. *Ecology* 57: 1104-1124

Breckle SW, Agachanjanz O, Rahmann M (1994) Spezielle Ökologie der gemäßigten und arktischen Zonen Euro-Nordasiens. Ed 2nd Vol Band 3. Gustav-Fischer Verlag. Stuttgart. Jena

Breman H, Kessler JJ (1995) Woody plants in agro-ecosystems of semi-arid regions. Springer-Verlag. Berlin

Briones O, Montaña C, Ezcurra E (1996) Competition between three Chihuahuan desert species: evidence from plant size-distance relations and root distribution. *Journal of Vegetation Science* 7: 453-460

Brisson J, Reynolds JF (1994) The effect of neighbors on root distribution in a creosotebush (*Larrea tridentata*) population. *Ecology* 75: 1693-1702

Brown JH, Jr, and Woods FW (1968) Root extension of trees in surface soils of the North Carolina piedmont. *Bot. Gaz.* 129: 1126-1132

Brown K (1992) *Prosopis cineraria* woodlands in Oman. past. present and future. In RW Dutton. M Powell. RJ Ridley. eds. *Prosopis species. aspects of their value. research and development*. Food and Agriculture Organisation of the United Nations. pp 131-144

Buettner V (1994) Spatial and temporal patterns of fine root abundance in a mixed oak-beech forest. *Forest ecology and management* 70:11–21.

Bunger MT, Thomson HJ (1938) Root development as a factor in the success or failure of windbreak trees in the southern High Plains. *Journal of Forestry* 36: 790-803

Burbridge NT (1945) Morphology and anatomy of the Western Australian species of *Triodia* R. Br. *Transactions of the Royal Society of South Australia* 69: 303-308

Burger JC, Louda SM (1995) Interaction of diffuse competition and insect herbivory in limiting brittle prickly pear cactus. *Opuntia fragilis* (Cactaceae). American Journal of Botany 82: 1558-1566

Burgess TL (1995) Desert grassland. mixed shrub savanna. shrub steppe. or semidesert scrub? The dilemma of coexisting growth forms. In MP McClaran. TR Van Devender. eds. *The desert grassland*. University of Arizona Press. Tucson. pp 31-67

Bursova TL (1983) Ekomorfoz kornevoj sistemy rastenij solonchakovych pochv drevnej Del'ty Cyrdar'i. (Ecomorphosis of plant root systems in the solonchak soils of the ancient Syr Darya delta (Kazakh SSR. USSR). In Russian). Izvestiia Akademii nauk Kazakhskoi SSR. Seriia biologicheskaya 1983: 6-13

Cable DR (1969) Competition in the semidesert grass-shrub type as influenced by root systems. growth habits. and soil moisture extraction. Ecology 50: 27-38

Campion WE (1926) The depth attained by roots. Australian Forestry Journal 9: 128

Canadell J, Jackson RB, Ehleringer JR, Mooney HA, Sala OE, Schulze ED (1996) Maximum rooting depth of vegetation types at the global scale. Oecologia 108: 583-595

Cannon HL (1960) The development of botanical methods of prospecting for uranium on the Colorado Plateau. In U.S. Geological Survey Bulletin. Vol 1085-A. United States Government Printing Office. Washington. D.C.. pp 1-50

Cannon WA (1911) The root habits of desert plants. Vol No. 131. Carnegie Institution of Washington. Washington. D.C.

Cannon WA (1913) Notes on root variation in some desert plants. The Plant World 16: 323-341

Cannon WA (1914) Specialization in vegetation and in environment in California. Plant World 17: 223-243

Cannon WA (1921) Plant habits and habitats in the arid portions of South Australia. Vol No. 308. Carnegie Institution of Washington. Washington. D.C.

Cannon WA (1924) General and physiological features of the vegetation of the more arid portions of southern Africa. with notes on the climatic environment. Vol No. 354. Carnegie Institution of Washington. Washington. D.C.

Carbon BA, Bartle GA, Murray AM, MacPherson DK (1980) The distribution of root length. and the limits to flow of soil water to roots in a dry sclerophyll forest. Forest Science 26: 656-664

Carter MR, Gregorich EG (2010) Carbon and nitrogen storage by deep-rooted tall fescue (*Lolium arundinaceum*) in the surface and subsurface soil of a fine sandy loam in eastern Canada. Agriculture, Ecosystems and Environment 136(1-2):125-132.

Clark RB, Alberts EE, Zoel RW, inclair TR, Miller MS, Kemper WD, Foy CD (1998) Eastern gamagrass (*Tripsacum dactyloides*) root penetration into and chemical properties of claypan soils. Plant and Soil 200: 33-45

Cody ML (1986) Structural niches in plant communities. In J Diamond. TJ Case. eds. *Community ecology*. Harper & Row Publishers. New York. pp 381-405

Coetzee JA, Page MI, Meredith D (1946) Root studies in Highveld grassland communities. South African Journal of Science 42: 105-118

Cole HE, Holch AE (1941) The root habits of certain weeds of southeastern Nebraska. *Ecology* 22: 141-147

Cottle HJ (1931) Studies in the vegetation of southwestern Texas. *Ecology* 12: 105-155

Cougenour MB, Ellis JE, Popp RG (1990) Morphometric relationships and developmental patterns of *Acacia tortilis* and *Acacia reficiens* in Southern Turkana. Kenya. *Bulletin of the Torrey Botanical Club* 117: 8-17

Coupland RT, Johnson RE (1965) Rooting characteristics of native grassland species in Saskatchewan. *Journal of Ecology* 53: 475-507

Currie PO, Hammer FL (1979) Detecting depth and lateral spread of roots of native range plants using radioactive phosphorus. *Journal of Range Management* 32: 101-103

Dabadghao PM, Marwaha SP, Gupta BS, Das RB, Deb Roy R (1963) Root ecology of some promising desert grasses of Rajasthan. *Annals of Arid Zone* 1: 163-173

Damman AWH (1971) Effect of vegetation changes on the fertility of a Newfoundland forest site. *Ecological Monographs* 41: 253-270

Das, DK, Chaturvedi OP (2008). Root biomass and distribution of five agroforestry tree species. *Agroforestry systems*, 74(3), 223.

Daubenmire RF (1941) Some ecologic features of the subterranean organs of alpine plants. *Ecology* 22(4):370–378.

David TS, et al. (2013) Root functioning, tree water use and hydraulic redistribution in *Quercus suber* trees: A modeling approach based on root sap flow. *Forest Ecology and Management* 307:136–146.

David TS, Ferreira MI, Cohen S, Pereira JS, David JS (2004) Constraints on transpiration from an evergreen oak tree in southern Portugal. *Agricultural and Forest Meteorology* 122(3–4):193–205.

Davidson E, Lefebvre PA, Brando PM, Ray DM, Trumbore SE, Solorzano LA, Nepstad D.C (2011) Carbon inputs and water uptake in deep soils of an eastern Amazon forest. *Forest Science*, 57(1), 51-58.

Davis CB (1972) Comparative ecology of six members of the *Arctostaphylos andersonii* complex. Ph.D. dissertation. University of California Davis. Davis

Davis EA, Pase CP (1977) Root system of shrub live oak: Implications for water yield in Arizona chaparral. *Journal of Soil and Water Conservation* 32: 174-180

Davis G, Neilsen W, Mcdavitt J (1983) Root distribution of *Pinus radiata* related to soil characteristics in five Tasmanian soils. *Australian Journal of Soil Research* 21(2):165.

Dawson T (1993) Hydraulic lift and water use by plants: Implications for water balance, performance and plant-plant interactions. *Oecologia* 95(4):565–574.

Day MW (1941) The root system of red pine saplings. *Journal of Forestry*:468–472.

Day MW. (1944). The root system of aspen. *American Midland Naturalist*, 32(2), 502.

Day WR (1959) Observations on eucalypts in Cyprus. II. Root development in relation to soil conditions. *Empire Forest Review* 38: 186-197

- Dell B, Bartle JR, Tacey WH (1983) Root occupation and root channels of jarrah forest subsoils. *Australian Journal of Botany* 31: 615–627
- Derbel S, Chaieb M (2012) Growth establishment and phenology of four woody Saharan species. *African Journal of Ecology* 51(2):307–318.
- DeSouza J, Silka PA, Davis SD (1986) Comparative physiology of burned and unburned *Rhus laurina* after chaparral wildfire. *Oecologia* 71: 63–68
- Dhyani SK, Narain P, Singh RK (1990) Studies on root distributions of five multipurpose tree species in Doon Valley. India. *Agroforestry Systems* 12: 149–161
- Dittmer HJ (1959) A study of the root system of certain sand dune plants in New Mexico. *Ecology* 40(2):265–273.
- Do FC, et al. (2008) Stable annual pattern of water use by *Acacia tortilis* in Sahelian Africa. *Tree physiology* 28(1):95–104.
- Dobrowolski JP, Caldwell MM, Richards JH (1990) Basin hydrology and plant root systems. In CB Osmond. LF Pitelka. GM Hidy. eds. *Plant biology of the basin and range*. Vol 80. Springer-Verlag. Berlin. pp 243–292
- Dodd J, Heddle EM, Pate JS, Dixon KW (1984) Rooting patterns of sandplain plants and their functional significance. In JS Pate. JS Beard. eds. *Kwongan: plant life on the sandplain*. University of Western Australia. Nedlands. pp 146–177
- Donovan LA, Richards JH, Muller MW (1996) Water relations and leaf chemistry of *Chrysothamnus nauseosus* ssp. *consimilis* (Asteraceae) and *Sarcobatus vermiculatus* (Chenopodiaceae). *American Journal of Botany* 83(12):1637–1646
- Dorji T, et al. (2013) Plant functional traits mediate reproductive phenology and success in response to experimental warming and snow addition in Tibet. *Global Change Biology* 19(2):459–472.
- Dougherty RL, Lauenroth WK, Singh JS (1996) Responses of a grassland cactus to frequency and size of rainfall events in a North American shortgrass steppe. *Journal of Ecology* 84: 177–183
- Douglas DA (1989) Clonal growth of *Salix setchelliana* on glacial river gravel bars in Alaska. *Journal of Ecology* 77(1):112–126.
- Drexhage M, Gruber F (1998) Architecture of the skeletal root system of 40-year-old *Picea abies* on strongly acidified soils in the Harz Mountains (Germany). *Canadian Journal of Forest Research* 28(1):13–22.
- Du J, et al. (2010) Clonal integration increases performance of ramets of the fern *Diplopterygium glaucum* in an evergreen forest in southeastern China. *Flora: Morphology, Distribution, Functional Ecology of Plants* 205(6):399–403.
- Dumortier M (1991) Below-ground dynamics in a wet grassland ecosystem. *Plant Root Growth, an Ecological Perspective*, ed Atkinson D (Blackwell Scientific Publications, Oxford).
- Duncan WH (1935) Root systems of woody plants of old fields of Indiana. *Ecology* 16(4):554–567.
- Dupuy NC, Dreyfus BL (1992) Bradyrhizobium populations occur in deep soil under the leguminous tree *Acacia albida*. *Applied and Environmental Microbiology* 58(8):2415–2419.

- Dye PJ (1996) Response of *Eucalyptus grandis* trees to soil water deficits. *Tree physiology* 16:233–238.
- Eamus D, Chen X, Kelley G, Hutley LB (2002) Root biomass and root fractal analyses of an open *Eucalyptus* forest in a savanna of north Australia. *Australian Journal of Botany* 50:31– 41.
- Eis S (1974) Root system morphology of western hemlock, western red cedar, and Douglas fir. *Canadian Journal of Forest Research* 4(1):28–38.
- Eis S (1987) Root systems of older immature hemlock, cedar, and Douglas-fir. *Canadian Journal of Forest Research* 17(11):1348–1354.
- Elliott GRB (1924) Relation between the downward penetration of corn roots and water level in peat soil. *Ecology* 5(2):175–178.
- Ellsworth PZ, Sternberg LSL (2015) Seasonal water use by deciduous and evergreen woody species in a scrub community is based on water availability and root distribution. *Ecohydrology* 5:51.
- Esler KJ, Rundel PW (1999) Comparative patterns of phenology and growth from diversity in two winter rainfall deserts: the Succulent Karoo and Mojave Desert ecosystems. *Plant Ecology* 142: 97-104
- Estrada-Medina H, Graham RC, Allen MF, Jimenez-Osornio JJ, Robles-Casolco S (2013) The importance of limestone bedrock and dissolution karst features on tree root distribution in northern Yucatan, Mexico. *Plant and Soil* 362(1–2):37–50.
- Evenari M (1938) Root conditions of certain plants of the Wilderness of Judaea. *Journal of the Linnean Society of London. Botany* 51: 383-388
- Evenari M, Shanan L, Tadmor N (1982) *The Negev: the challenge of a desert*. Ed 2nd. Harvard University Press. Cambridge. Massachusetts
- Fagg C (1991) *Acacia tortilis: Fodder tree for desert sands*. Vol 91. Nitrogen Fixing Tree Association. Hawaii.
- Fan Y, et al. (2012) Water adaptive traits of deep-rooted C3 halophyte (*Karelinia caspica* (Pall.) Less.) and shallow-rooted C4 halophyte (*Atriplex tatarica* L.) in an arid region, Northwest China. *Journal of Arid Land* 4(4):469–478.
- Fan Y, Miguez-Macho G, Jobbágy EG, Jackson RB, & Otero-Casal C (2017). Hydrologic regulation of plant rooting depth. *Proceedings of the National Academy of Sciences*, 114(40), 10572-10577.
- Farrington P, Greenwood EAN, Bartle GA, Beresford JD, Watson GD (1989) Evaporation from Banksia woodland on a groundwater mound. *Journal of Hydrology* 105(1–2):173–186.
- Farrish KW (1991) Spatial and temporal fine-root distribution in three Louisiana forest soils. *Soil Science Society of America journal* 55(6):1752–1757.
- Fayle DCF (1965). Rooting habit of sugar maple and yellow birch. Published under the authority of the Minister of Forestry.
- February EC, Allsopp N, Shabane T, Hattas D (2011) Coexistence of a C4 grass and a leaf succulent shrub in an arid ecosystem. The relationship between rooting depth, water and nitrogen. *Plant and Soil* 349(1–2):253–260.
- February EC, Cook GD, Richards AE (2013) Root dynamics influence tree-grass coexistence in an Australian savanna. *Austral Ecology* 38(1):66–75.

- Fenner M (1980) Some measurements on the water relations of Baobab trees. *Biotropica* 12(3):205–209.
- Fernández RJ, Paruelo JM (1988) Root systems of two Patagonian shrubs: A quantitative description using a geometrical method. *Journal of Range Management* 41: 220-223
- Fisher RA, et al. (2007) The response of an Eastern Amazonian rain forest to drought stress: results and modelling analyses from a throughfall exclusion experiment. *Global Change Biology* 13(11):2361–2378.
- Flombaum P, Sala O (2011) Effects of plant species traits on ecosystem processes. *Ecology* 93(2):227–234.
- Foldats, Ernesto, Rutkis E (1975) Ecological studies of Chaparro (*Curatella americana* L.) and Manteco (*Byrsonima crassifolia* H.B.K.) in Venezuela. *Ecological Studies* 2(3):159– 178.
- Follett R, Allmaras R, Reichman G (1974) Distribution of corn roots in sandy soil with a declining water table. *AGRONOMY JOURNAL* 66:288–292.
- Fonteyn PJ, Mahall BE (1981) An experimental analysis of structure in a desert plant community. *Journal of Ecology* 69: 883-896
- Forseth IN, Ehleringer JR, Werk KS, Cook CS (1984) Field water relations of Sonoran Desert annuals. *Ecology* 65: 1436-1444
- Frangi JL, Lugo AE (1985) Ecosystem dynamics of a subtropical floodplain forest. *Ecological Monographs* 55(3):351–369.
- Franzluebbers AJ, Stuedemann JA (2009) Soil-profile organic carbon and total nitrogen during 12 years of pasture management in the Southern Piedmont USA. *Agriculture, Ecosystems and Environment* 129(1–3):28–36.
- Freckman DW, Virginia RA (1989) Plant-feeding nematodes in deep-rooting desert ecosystems. *Ecology* 70: 1665-1678
- Freycon V, et al. (2014) Tree roots can penetrate deeply in African semi-deciduous rain forests: evidence from two common soil types. *Journal of Tropical Ecology* 31(1):13–23.
- Frith AC (1955) No man's land. *Empire Forestry Review* 34: 179-187
- Gaines KP, et al. (2016) Reliance on shallow soil water in a mixed-hardwood forest in central Pennsylvania. *Tree Physiology* 36(4):444–458.
- Gary HL (1963) Root distribution of five-stamen tamarisk. seepwillow. and arrowweed. *Forest Science* 9: 311-314
- Gaze SR, Brouwer J, Simmonds LP, Bromley J (1998) Dry season water use patterns under *Guiera senegalensis* L. shrubs in a tropical savanna. *Journal of Arid Environments* 40: 53-67
- Gemmer EW (1928) The root system of a longleaf pine. *The Scientific Monthly* 27(4):384–384.
- Gentile RM, Martino DL, Entz MH (2003) Root characterization of three forage species grown in southwestern Uruguay. *Canadian Journal of Plant Science* 83(4):785–788.
- George AS, Hopkins AJM, Marchant NG (1979) The heathlands of western Australia. In RL Specht. ed. *Heathlands and related shrublands*. Vol 9A. Elsevier. Amsterdam. pp 211-230

Germon A, et al. (2016) Unexpected phenology and lifespan of shallow and deep fine roots of walnut trees grown in a silvoarable Mediterranean agroforestry system. *Plant and Soil* 401(1–2):409–426.

Gibbens RP, Lenz JM (2001) Root systems of some Chihuahuan Desert plants. *Journal of Arid Environments*

Gile LH, Gibbens RP, Lenz JM (1995) Soils and sediments associated with remarkable, deeply-penetrating roots of crucifixion thorn (*Koeberlinia spinosa* Zucc.). *Journal of Arid Environments* 31: 137-151

Gile LH, Gibbens RP, Lenz JM (1998) Soil-induced variability in root systems of creosotebush (*Larrea tridentata*) and tarbush (*Flourensia cernua*). *Journal of Arid Environments* 39: 57-78

Giliberto J, Estay H (1978) Seasonal water stress in some Chilean matorral shrubs. *Botanical Gazette* 139: 236-240

Ginzburg C (1966) Xerophytic structures in the roots of desert shrubs. *Annals of Botany* 30: 403-418

Glover JD, et al. (2010) Harvested perennial grasslands provide ecological benchmarks for agricultural sustainability. *Agriculture, Ecosystems and Environment* 137(1–2):3–12.

Glover PE (1950) The root systems of some British Somaliland plants--I. *East African Agricultural Journal* 16: 98-113

Glover PE (1951) The root systems of some British Somaliland plants--II. *East African Agricultural Journal* 16: 154-162

Glover PE (1951) The root systems of some British Somaliland plants--III. *East African Agricultural Journal* 16: 205-217

Glover PE (1951) The root systems of some British Somaliland plants--IV. *East African Agricultural Journal* 17: 38-50

Glover PE, Trump EC, Waterride LED (1964) Termitaria and vegetation patterns on the Loita Plains of Kenya. *Journal of Ecology* 52: 367-377

Golluscio RA, Sala OE (1993) Plant functional types and ecological strategies in Patagonian forbs. *Journal of Vegetation Science* 4: 839-846

Golovina RD (1991) Root systems of juniper forest of the Alai Ridge. *Soviet Forest Sciences* 1991: 64-67

Goor AY, Barney CW (1968) Forest tree planting in the arid zones. The Ronald Press Company. New York

Greenland DJ, Kowal JML (1960) Nutrient content of the moist tropical forest of Ghana. *Plant and Soil* 12(2):154–174.

Gries D, et al. (2003) Growth and water relations of *Tamarix ramosissima* and *Populus euphratica* on Taklamakan desert dunes in relation to depth to a permanent water table. *Plant, Cell and Environment* 26(5):725–736.

Groeneveld DP (1989) Shrub rooting and water acquisition on threatened shallow groundwater habitats in the Owens Valley, California. In *Proceedings--Symposium on cheatgrass invasion, shrub die-off, and other aspects of shrub biology and management*. Vol INT-276. U.S.D.A. Forest Service Intermountain Research Station. Ogden, Utah. pp 221-237

- Groeneveld DP, Crowley DE (1988) Root system response to flooding in three desert shrub species. *Functional Ecology* 2(4):491.
- Groot JJR, Koné D, Traoré M, Kamissoko N (1995) Description du système racinaire de l'*Andropogon gayanus*, du *Vigna unguiculata* et du *Stylosanthes hamata* en zone soudano-sahélienne. Vol 8. Production Soudano-Sahélienne (PSS). Wageningen
- Guinness (2003) Guinness world records 2004. Guinness World Records LTD.. New York
- Gulmon SL, Mooney HA (1979) Spatial relationships and competition in a Chilean desert cactus. *Oecologia* 44: 40-43
- Haag CL, Johnson JE, Erdmann GG (1989) Rooting depths of red maple (*Acer rubrum* L.) on various sites in the Lake States. US Dep. Agric. For. Serv. Res. Note NC-347.
- Haase P, et al. (1996) An investigation of rooting depth of the semiarid shrub *Retamasphaerocarpa* (L.) Boiss. by labelling of ground water with a chemical tracer. *Journal of Hydrology* 177:23–31.
- Haasis FW (1921) Relations between soil type and root form of western yellow pine seedlings. *Ecology* 2(4):292–303.
- Haigh H (1966) Root development in the sandy soils of Zulu land. *Forestry in South Africa* 7:31–36.
- Hanes TL (1965) Ecological studies on two closely related chaparral shrubs in southern California. *Ecological Monographs* 35: 213-235
- Hao Y, et al. (2006) Roots of pioneer trees in the lower sub-tropical area of Dinghushan, Guangdong, China. *Journal of Zhejiang University Science B* 7(5):377–385.
- Harris GA (1967) Some competitive relationships between *Agropyron spicatum* and *Bromus tectorum*. *Ecological Monographs* 37: 89-111
- Haworth F (1953) Observations on the root system of Guatemala grass on an upland tea soil in Ceylon. *Tropical Agriculture* 30: 116-121
- Hayes FA, Stoeckeler JH (1935) Soil and forest relationships of the shelter beltzone. In USFS The Lake States Forest Experiment Station. ed. Possibilities of shelterbelt planting in the Plains Region. United States Government Printing Office. Washington. D.C.. pp 111-153
- He WM, Zhang XS (2003) Responses of an evergreen shrub *Sabina vulgaris* to soil water and nutrient shortages in the semi-arid Mu Us Sandland in China. *Journal of Arid Environments* 53(3):307–316.
- Hellmers H, Horton JS, Juhren G, O'Keefe J (1955) Root systems of some chaparral plants in southern California. *Ecology* 36: 667-678
- Hendrick RL, Pregitzer KS, Hendrick RL, Pregitzer KS (1996) Temporal and Depth-Related Patterns of Fine Root Dynamics in Northern Hardwood Forests. *Journal of Ecology* 84(2):167–176.
- Heyward F (1933) The root system of longleaf pine on the deep sands of western Florida. *Ecology* 14: 136-148
- Higgins KB, Lamb AJ, van Wilgen BW (1987) Root systems of selected plant species in mesic mountain fynbos in the Jonkershoek Valley. south-western Cape Province. *South African Journal of Botany* 53: 249-257

Hipondoka MHT, Aranibar JN, Chirara C, Lihavha M, Macko SA (2003) Vertical distribution of grass and tree roots in arid ecosystems of Southern Africa: niche differentiation or competition? *Journal of Arid Environments* 54(2):319–325.

Hironaka M (1961) The Relative Rate of Root Development of Cheatgrass and Medusahead. *Journal of Range Management Archives* 14(5):263–267.

Hnatiuk RJ, Hopkins AJM (1980) Western Australian species-rich kwongan (sclerophyllous shrubland) affected by drought. *Australian Journal of Botany* 28: 573-585

Hoffmann A (1978) Root studies in the Chilean matorral. *Oecologia* 1 32(1):57–69.

Holdo RM, Timberlake J (2008) Rooting depth and above-ground community composition in Kalahari sand woodlands in western Zimbabwe. *Journal of Tropical Ecology* 24(2):169–176.

Hopkins HH (1951) Ecology of the native vegetation of the loess hills in central Nebraska. *Ecological Monographs* 21: 125-147

Horton KW (1958) Rooting habits of lodgepole pine doi:10.1038/180270a0.

Hosegood PH (1963) The root distribution of Kikuyu grass and Wattle trees. *East African Agricultural and Forestry Journal* 29: 60-61

Hosegood PH, Howland P (1966) A preliminary study of the root distribution of some exotic tree crops, evaluated by a rapid sampling method. *East African agricultural and forestry journal* 32:16–18.

Howard A (1925) The effect of grass on trees. *Proceedings of the Royal Society of London. Series B* 97: 284-321

Hubble TCT, Docker BB, Rutherford ID (2010) The role of riparian trees in maintaining riverbank stability: A review of Australian experience and practice. *Ecological Engineering* 36(3):292–304.

Hulbert LC (1955) Ecological studies of Bromus tectorum and other annual bromegrasses. *Ecological Monographs* 25: 181-213

Hull JC, Muller CH (1977) The potential for dominance by Stipa pulchra in a California grassland. *American Midland Naturalist* 97: 147-175

Humphrey RR (1937) Ecology of the burroweed. *Ecology* 18: 1-9

Ignatenko IV, Knorre AV, Lovelius NV, Norin BN (1972) Standing crop in plant communities at the station Ary-Mas. In FE Wielgolaski. T Rosswall. eds. *Tundra biome*. Tundra Biome Steering Committee. Stockholm. pp 140-148

Imai N, Kitayama K, Titin J (2010) Distribution of phosphorus in an above-to-below-ground profile in a Bornean tropical rain forest. *Journal of Tropical Ecology* 26(6):627–636.

Jackson R, Banner J, Jobbágy E (2002) Ecosystem carbon loss with woody plant invasion of grasslands. *Nature* 277(July):623–627.

Jackson RB, Moore LA, Hoffmann WA, Pockman WT, Linder CR (1999) Ecosystem rooting depth determined with caves and DNA. *Proceedings of the National Academy of Sciences (USA)* 96: 11387-11392

Janzen DH (1970) *Jaqinina pungens*. an heliophile from the understory of tropical deciduous forests. *Biotropica* 2: 112-119

Jaramillo VJ, Ahedo-Hernández R, Kauffman JB (2003) Root biomass and carbon in a tropical evergreen forest of Mexico: changes with secondary succession and forest conversion to pasture. *Journal of Tropical Ecology* 19(4):457–464.

Jeffers JNR, Boaler SB (1966) Ecology of a Miombo site. Lupa North Forest Reserve. Tanzania I. Weather and plant growth. 1962-64. *Journal of Ecology* 54: 447-463

Jeník J (1971) Root structure and underground biomass in equatorial forests. In P Duvigneaud. ed. *Productivity of forest ecosystems*. Vol 4. Unesco. Paris. pp 323-331

Jennings CMH (1974) The hydrology of Botswana. Ph.D. thesis. University of Natal. Pietermaritzburg. South Africa

Jiménez EM, Moreno FH, Lloyd J, Peñuela MC, Patiño S (2009) Fine root dynamics for forests on contrasting soils in the Colombian Amazon. *Biogeosciences Discussions* 6(2):3415–3453.

Jipp PH, Nepstad DC, Cassel DK, Reis De Carvalho C (1998) Deep soil moisture storage and transpiration in forests and pastures of seasonally-dry Amazonia. *Climatic Change* 39:395– 412.

Jobbagy EG, Jackson RB (2004) Groundwater use and salinization with grassland afforestation. *Global Change Biology* 10(8):1299–1312.

Jobbágy EG, Nusetto MD, Villagra PE, Jackson RB (2011) Water subsidies from mountains to deserts: Their role in sustaining groundwater-fed oases in a sandy landscape. *Ecological Applications* 21(3):678–694.

Joffre R, Leiva Morales MJ, Rambal S, Fernández Ales R (1987) Dynamique racinaire et extraction de l'eau du sol par des graminées pérennes et annuelles méditerranéennes. *Acta Oecologica. Oecologia Plantarum* 8: 181-194

Johnson DM, Domec JC, Woodruff DR, McCulloh KA, Meinzer FC (2013) Contrasting hydraulic strategies in two tropical lianas and their host trees. *American Journal of Botany* 100(2):374–383.

Johnson KL, Wright GM, Ashton DH (1968) Ecological studies of Tunnel Cave. Mt. Eccles. Victoria Naturalist 85: 350-356

Jones RH, Lockaby BG, Somers GL (1996) Effects of microtopography and disturbance on fine-root dynamics in wetland forests of low-order stream floodplains. *American Midland Naturalist* 136(1):57–71.

Jonsson K, Fidjeland L, Maghembe JA, Hogberg P (1988) The vertical distribution of fine roots of five tree species and maize in Morogoro, Tanzania. *Agroforestry Systems* 6:63–69.

Joslin JD, Gaudinski JB, Torn MS, Riley WJ, Hanson PJ (2006) Fine-root turnover patterns and their relationship to root diameter and soil depth in a 14C-labeled hardwood forest. *New Phytologist* 172(3):523–535.

Kalisz P, Zimmerman RW, Muller RN (1987) Root density, abundance, and distribution in the mixed mesophytic forest of eastern Kentucky. *Soil Science Society of America journal* 51(1):220–225.

Kalisz, PJ, Stringer JW, Volpe JA, Clark. DT (1988) Trees as monitors of tritium in soil water. *J. Environ. Qual.*. 17: 62-70

Karasz I (1996) The root system of *Juniperus communis* L. in a sandy soil in Central Hungary. *Acta Phytogeogr Suec* 81:32–34.

Karizumi N (1979) Monograph - Root system forms and distributions of individual trees in Japan (in Japanese) (Seibundo Shinkosha Publishing Co. Ltd., Tokyo) Available at:<https://www.cabdirect.org/cabdirect/abstract/19870618641>.

Karpov VG, ed (1983) Faktory reguliatsii ekosistem elovykh lesov (Regulation factors of spruce forest ecosystems. In Russian). Nauka. Leningrad

Karrfalt EE (1981) The comparative and developmental morphology of the root system of *Selaginella selaginoides* (L.) Link. *American journal of botany* 68(2):244–253.

Kausch W (1960) Der Einfluß von edaphischen und klimatischen Faktoren auf die Ausbildung des Wurzelwerkes der Pflanzen unter besonderer Berücksichtigung einiger algerischer Wüstenpflanzen. Habilitation. Technische Hochschule Darmstadt. Darmstadt

Keller B (1930) Die Methoden zur Erforschung der Ökologie der Steppen- und Wüstenpflanzen. In E Abderhalden. ed. *Handbuch der biologischen Arbeitsmethoden*. Vol 11(6). Urban & Schwarzenberg. Berlin. pp 1-128

Kellman M (1990) Root proliferation in recent and weathered sandy soils from Veracruz, Mexico. *Journal of Tropical Ecology* 6(3):355.

Kenzo T, et al. (2009) Development of allometric relationships for accurate estimation of above- and below-ground biomass in tropical secondary forests in Sarawak, Malaysia. *Journal of Tropical Ecology* 25(4):371.

Keresztesi B (1968) Morphological characteristic of the *Robinia* root system on different sites of the Great Hungarian Plain. In MS Ghilarov. VA Kovda. LN Novichkova-Ivanova. LE Rodin. VM Sveshnikova. eds. *Methods of productivity studies in root systems and rhizosphere organisms*. International Symposium USSR. August 28–September 12. 1968. Nauka. Leningrad. pp 86-96

Kerfoot O (1963) The root systems of tropical forest trees. *Commonwealth Forestry Review* 42: 19-26

Kimber PC (1974) The root system of jarrah (*Eucalyptus marginata*). Vol 10. Western Australia Forestry Department. Perth. Australia

Kleinhampl FJ, Koteff C (1960) Botanical Prospecting for U rani urn in the Circle Cliffs Area Garfield County , Utah, USGS Bulletin 1085-C (USGS).

Klepper EL, Gano KA, Cadwell LL (1985) Rooting depth and distributions of deep-rooted plants in the 200 Area Conrol Zone of the Hanford site. Vol PNL-5247. Pacific Northwest Laboratory. Richland. Washington

Kohzu A, Matsui K, Yamada T, Sugimoto A, Fujita N (2003) Significance of rooting depth in mire plants: Evidence from natural ^{15}N abundance. *Ecological Research* 18(3):257–266.

Kokoreva II (1996) Root systems of *Cratae gus* L. in the Trans-Ili Alatau, Kazakhstan. *Acta Phytogeogr Suec* 81:35–38.

Kong Lingshao MM (1996) Reserach on bio-ecological characteristics of *Suaeda physophora* and its community at the Hutubi cattle farm of Xinjiang. China. *Acta Botanica Sinica* 38: 475-482

Köstler JN, Brückner E, Bibelriether H (1968) Die Wurzeln der Waldbäume: Untersuchungen zur Morphologie der Waldbäume in Mitteleuropa. Paul Parey. Hamburg. Berlin

Koteen LE, Baldocchi DD, Harte J (2011) Invasion of non-native grasses causes a drop in soil carbon storage in California grasslands. *Environmental Research Letters* 6:44001.

Kreutzer K (1968) The root system of the red alder (*Alnus glutinosa* Gärtn.). In MS Ghilarov. VA Kovda. LN Novichkova-Ivanova. LE Rodin. VM Sveshnikova. eds. Methods of productivity studies in root systems and rhizosphere organisms. International Symposium USSR. August 28--September 12. 1968. Nauka. Leningrad. pp 114-119

Krishnamurthy L, Zaman-Allah M, Marimuthu S, Wani SP, Kesava Rao AVR (2012) Root growth in *Jatropha* and its implications for drought adaptation. *Biomass and Bioenergy* 39:247–252.

Kubota M, Tenhunen J, Zimmermann R, Adiku S, Kakubari Y (2005) Influences of environmental factors on the radial profile of sap flux density in. *Tree Physiology* (1998):545–556.

Kuiper LC (1992) Spatial disposition and extension of the structural root system of Douglas-fir. *Forest ecology and management* 47(1):111–125.

Kummerow J (1981) Structure of roots and root systems. In F DiCastri. DW Goodall. RL Specht. eds. Mediterranean-type shrublands. Elsevier. Amsterdam. pp 269-288

Kutschera L, Lichtenegger E (1960) Wurzelatlas mitteleuropäischer Ackerunkräuter und Kulturpflanzen. DLG-Verlags GmbH. Frankfurt am Main

Kutschera L, Lichtenegger E, Sobotik M (1982) Wurzelatlas mitteleuropäischer Grünlandpflanzen. Band 1: Monocotyledonae. Gustav Fischer. Stuttgart

Kutschera L, Lichtenegger E, Sobotik M (1992) Wurzelatlas mitteleuropäischer Grünlandpflanzen. Band 2: Pteridophyta und Dicotyledonae (Magnoliopsida). Teil 1: Morphologie. Anatomie. Ökologie. Verbreitung. Soziologie. Wirtschaft. Gustav Fischer. Stuttgart

Kutschera L, Lichtenegger E, Sobotik M (1997a) Bewurzelung von Pflanzen in verschiedenen Lebensräumen. Vol 49. Land Oberösterreich. OÖ. Landesmuseum. Linz. Austria

Kutschera L, Lichtenegger E, Sobotik M (2009) Wurzelatlas der Kulturpflanzen gemäßiger Gebiete mit Arten des Feldgemüsebaues. DLG-Verlag Frankfurt am Main ISBN-Nr.: 978-3-7690-0708-4

Kutschera L, Lichtenegger E, Sobotik M, Haas D (1997b) Die Wurzel. das neue Organ. ihre Bedeutung für das Leben von *Welwitschia mirabilis* und anderer Arten der Namib. sowie von Arten angrenzender Gebiete. Pflanzensoziologisches Institut. Klagenfurt. Austria

Kutschera L, Lichtenegger E (2002). Wurzelatlas mitteleuropäischer Waldbäume und Sträucher. Leopold Stocker Verlag Graz und Stuttgart. ISBN-Nr.: 978-3-7020-0928-0

Laclau JP, et al. (2004) The function of the superficial root mat in the biogeochemical cycles of nutrients in congolesse *Eucalyptus* plantations. *Annals of Botany* 93(3):249–261.

Laclau JP, et al. (2013) Dynamics of soil exploration by fine roots down to a depth of 10 m throughout the entire rotation in *Eucalyptus grandis* plantations. *Frontiers in plant science* 4(July):243.

Laclau JP, Arnaud M, Bouillet JP, Ranger J (2001) Spatial distribution of Eucalyptus roots in a deep sandy soil in the Congo: relationships with the ability of the stand to take up water and nutrients. *Tree Physiology* 21: 129–136.

Lamont BB, Lange BJ (1976) "Stalagmiform" roots in limestone caves. *New Phytologist* 76: 353-360

Leduc G, Favreau G, Schroeter P (2001) Long-term rise in a Sahelian water-table: the Continental Terminal in South-West Niger. *Journal of Hydrology* 24: 43-54

Lee CA, Lauenroth WK (1994) Spatial distribution of grass and shrub root systems in the shortgrass steppe. *American Midland Naturalist* 132: 117-123

Leistner OA (1967) The plant ecology of the southern Kalahari. *Botanical Survey of South Africa Memoirs* 38: 172

Lewis DC, Burgoyne RH (1964) The relationship between oak tree roots and groundwater in fractured rock as determined by tritium tracing. *Journal of Geophysical Research* 69: 2579-2588

Lichtenegger E (1996) Root distribution in some alpine plants. *Acta Phytogeographica Suecica* 81: 76-82

Link SO, Cadwell LL, Petersen KL, Sackschewsky MR, Landeen DS (1995) The role of plants and animals in isolation barriers at Hanford. Washington. Vol PNL-10788. Pacific Northwest Laboratory. Richland. Washington

Link SO, Gee GW, Downs JL (1990) The effect of water stress on phenological and ecophysiological characteristics of cheatgrass and Sandberg's bluegrass. *Journal of Range Management* 43: 506-513

Louw GN, Seely MK (1980) Exploitation of fog water by a perennial Namib dune grass. *Stipagrostis sabulicola*. *South African Journal of Science* 76: 38-39

Ludwig JA (1977) Distributional adaptations of root systems in desert environments. In JK Marshall, ed. *The belowground ecosystem: a synthesis of plant-associated processes*. Vol 26. Colorado State University. Fort Collins. Colorado. pp 85-91

Lyford WH (1980) Development of the root system of northern red oak (*Quercus rubra* L.) Harvard For. Bull. 21. 30 pp.

Manning SJ, Barbour MG (1988) Root systems, spatial patterns, and competition for soil moisture between two desert shrubs. *American Journal of Botany* 75: 885-893

Manning SJ, Groeneveld DP (1989) Shrub rooting characteristics and water acquisition on xeric sites in the western Great Basin. In *Proceedings--Symposium on cheatgrass invasion, shrub die-off, and other aspects of shrub biology and management*. Vol INT-276. U.S.D.A. Forest Service Intermountain Research Station. Ogden. Utah. pp 238-244

Markle MS (1917) Root systems of certain desert plants. *Botanical Gazette* 64: 177-205

McKell CM, Jones MB, Perrier ER (1962) Root production and accumulation of root material on fertilized annual range. *Agronomy Journal* 54: 459-462

Meinzer OE (1927) Plants as indicators of groundwater. Vol 577. Department of the Interior. U. S. Geological Survey. Washington. D.C.

Mensah KOA, Jenik J (1968) Root system of tropical trees 2. Features of the root system of iroko (*Chlorophora excelsa* Benth. et Hook.). *Preslia* 40: 21-27

Midwood AJ, Boutton TW, Archer SR, Watts SE (1998) Water use by woody plants on contrasting soils in a savanna parkland: assessment with d₂H and d₁₈O. *Plant and Soil* 205: 13-24

Migahid AM (1961) The drought resistance of Egyptian desert plants. *UNESCO Arid Zone Research* 16: 213-233

Miller PC, Ng E (1977) Root:shoot biomass ratios in shrubs in southern California and central Chile. *Madroño* 24: 215-223

Millikin CS, Bledsoe CS (1999) Biomass and distribution of fine and coarse roots from blue oak (*Quercus douglasii*) trees in the northern Sierra Nevada foothills of California. *Plant and Soil* 214: 27-38

Miroshnichenko YM (1975) Kornevye sistemy drevesnykh i kustarnikovych rastenij i ich zkologija v vostocnykh karakumach. (Root systems of trees and bushes and their ecology in eastern Karakum. In Russian). *Botanicheskii Zhurnal* 60: 1776-1795

Mooney HA, Gulmon SL, Rundel PW, Ehleringer JR (1980) Further observations on the water relations of *Prosopis tamarugo* of the northern Atacama Desert. *Oecologia* 44: 177-180

Mordelet P, Barot S, Abbadie L (1996) Root foraging strategies and soil patchiness in a humid savanna. *Plant and Soil* 182: 171-176

Moreira MZ, Sternberg LdSL, Nepstad D (2000) Vertical patterns of soil water uptake by plants in a primary forest and an abandoned pasture in the eastern Amazon: an isotopic approach. *Plant and Soil* 222: 95-107

Morello J (1955) Estudios botánicos en las regiones áridas de la Argentina. II. Transpiración de los arbustos resinosos de follaje permanente del Monte. *Revista Agronomica del Noroeste Argentino* 1: 385-524

Morello J (1956) Estudios botánicos en las regiones áridas de la Argentina. III. *Revista Agronomica del Noroeste Argentino* 2: 79-152

Morello J (1958) La provincia fitogeográfica del Monte. University of Tucuman. Tucuman. Argentina

Muller CH (1946) Root development and ecological relations of guayule. Vol No. 923. United States Department of Agriculture. Washington. D. C.

Nechaeva NT (1985) Description of plants used for vegetative range improvement. In NT Nechaeva. ed. Improvement of desert ranges in Soviet Central Asia. Vol 4. Harwood Academic Publishers. Chur. Switzerland. pp 55-120

Nechaeva NT, ed (1985) Improvement of desert ranges in Soviet Central Asia. Vol 4. Harwood Academic Publishers. Chur. Switzerland

Nechaeva NT. Prikhod'ko SY, eds (1968) Sown winter ranges in the foothill deserts of Soviet Central Asia. Israel Program for Scientific Translations. Jerusalem

Nesterova SG (1996) Root structure of plants in the Zailiisky Alatau Range. Kazakhstan. *Acta Phytogeographica Suecica* 81: 86-87

Nijland W, van der Meijde M, Addink EA, de Jong SM (2010) Detection of soil moisture and vegetation water abstraction in a mediterranean natural area using electrical resistivity tomography. *Catena* 81(3):209–216.

Nobel PS (1989) Temperature, water availability, and nutrient levels at various soil depths: Consequences for shallow-rooted desert succulents, including nurse plant effects. American Journal of Botany 76: 1486-1492

Nobel PS, Franco AC (1986) Annual root-growth and intraspecific competition for a desert bunchgrass. Journal of Ecology 74: 1119-1126

Nobel PS, Loik ME, Meyer RW (1991) Microhabitat and diel tissue acidity changes for two sympatric cactus species differing in growth habit. Journal of Ecology 79: 167-182

Nobel PS, Zutta BR (2005) Morphology, ecophysiology, and seedling establishment for *Fouquieria splendens* in the northwestern Sonoran Desert. J. Arid Environ. 62, 251-26510.1016/j.jaridenv.2004.11.002 (doi:10.1016/j.jaridenv.2004.11.002)

Olson DM, Dinerstein E, Wikramanayake ED, Burgess ND, Powell GVN, Underwood EC, D'amico JA, Itoua I, Strand HE, Morrison JC et al. 2001. Terrestrial ecoregions of the world: a new map of life on Earth: a new global map of terrestrial ecoregions provides an innovative tool for conserving biodiversity. BioScience 51: 933-938.

Osman A, Pieper RD (1988) Growth of *Gutierrezia sarothrae* seedlings in the field. Journal of Range Management 41: 92-93

Orcival JM, Floret C, Le Floc'h E, Ponantier R (1994) Water relations between two perennial species in the steppes of southern Tunisia. Journal of Arid Environments 28: 333-350

Pavlychenko TK (1937) Quantitative study of the entire root systems of weed and crop plants under field conditions. Ecology 18: 62-79

Peláez DV, Distel RA, Bóo RM, Elia OR, Mayor MD (1994) Water relations between shrubs and grasses in semi-arid Argentina. Journal of Arid Environments 27: 71-78

Phillips WS (1963) Depth of roots in soil. Ecology 44: 424

Popov KP (1979) Fistashka v srednei Azii. (The pistachio in central Asia. In Russian). Ylym. Ashkhabad

Preston CE (1900) Observations on the root system of certain Cactaceae. Botanical Gazette 30: 348-351

Ramam SS (1970) Root development in alluvial grasslands of Varanasi. Indian Forester 96: 100-110

Rawitscher F (1948) The water economy of the vegetation of the 'campos cerrados' in southern Brazil. Journal of Ecology 36: 237-268

Reynolds TD, Fraley L, Jr. (1989) Root profiles of some native and exotic plant species in southeastern Idaho. Environmental and Experimental Botany 29: 241-248

Roberts EA, Herty SD (1934) *Lycopodium Complanatum* Var.*Flabelliforme* Fernald: Its Anatomy And A Method Of Vegetative Propagation. American Journal of Botany 21.10:688-697. Web.

Roskov Y, Abucay L, Orrell T, Nicolson D, Bailly N, Kirk PM, Bourgoin T, DeWalt RE, Decock W, De Wever A et al. 2017. Species 2000 & ITIS catalogue of life, 2017 annual checklist. Species 2000. Leiden, the Netherlands: Naturalis. Digital resource at [WWW document] URL www.catalogueoflife.org/annual-checklist/2017 [accessed 1 September 2020].

Roupsard O, Ferhi A, Granier A, Pallo F, Depommier D, Mallet B, Joly HI, Dreyer E (1999) Reverse phenology and dry-season water uptake by *Faidherbia albida* (Del.) A. Chev. in an agroforestry parkland of Sudanese west Africa. *Functional Ecology* 13: 460-472

Roux J, Hopper S, Smith R (2009). *Isoetes eludens* (Isoetaceae), a New Endemic Species from the Kamiesberg, Northern Cape, South Africa. *Kew Bulletin*, 64(1), 123-128. Retrieved from <http://www.jstor.org/stable/20649631>

Rundel PW, Ehleringer JR, Mooney HA, Gulmon SL (1980) Patterns of drought response in leaf-succulent shrubs of the coastal Atacama Desert in northern Chile. *Oecologia* 46: 196-200

Rutherford MC (1983) Growth rates, biomass and distribution of selected woody plant roots in *Burkea africana*--*Ochna pulchra* savanna. *Vegetatio* 52: 45-63

Saunier RE Wagle RF (1967) Factors affecting the distribution of shrub live oak (*Quercus turbinella Greene*). *Ecology* 48: 35-41

Saurina NI Kamenetskaya IV (1969) Massa kornej sosny obyknovennoj (*Pinus sylvestris L.*) v sosnjake mshisto-lishajnikovom juzhnoj tajgi. (The root mass of *Pinus sylvestris L.* in moss-lichen pine forests of the southern taiga. In Russian with English summary). *Biulleten Moskovskogo obshchestva ispytatelei prirody. Otdel biologicheskii* 74: 96-100

Savory BM (1963) Site quality and tree root morphology in northern Rhodesia. *Rhodesian Journal of Agricultural Research* 1: 55-65

Schulze E-D, Mooney HA, Sala OE, Jobbág E, Buchmann N, Bauer G, Canadell J, Jackson RB, Loret J, Oesterheld M, Ehleringer JR (1996) Rooting depth, water availability, and vegetation cover along an aridity gradient in Patagonia. *Oecologia* 108: 503-511

Schuster JL (1964) Root development of native plants under three grazing intensities. *Ecology* 45: 63-70

Scott JD, van Breda NG (1937a) Preliminary studies on the root system of *Galenia africana* on the Worcester Veld Reserve. *South African Journal of Science* 34: 268-274

Scott JD, van Breda NG (1937b) Preliminary studies on the root system of the *Rhenosterbos* (*Elytropappus rhinocerotis*) on the Worcester Veld Reserve. *South African Journal of Science* 33: 560-569

Scott JD, van Breda NG (1939) Preliminary studies on the root system of *Euphorbia mauretanica*. *Euphorbia burmanni* and *Ruschia multiflora* on the Worcester Veld Reserve. *South African Journal of Science* 36: 227-235

Sen DN (1982) Environment and plant life in Indian Desert. Geobios International. Jodhpur. India

Sen DN, Tanwar GS (1983) Arid environment and root behaviour. In W Böhm, L Kutschera, E Lichtenegger, eds. *Wurzelökologie und ihre Nutzanwendung/Root ecology and its practical application*. Bundesanstalt für alpenländische Landwirtschaft Grumpenstein. Irdning. F. R. Germany. pp 185-206

Shachori A, Rosenzweig D, Poljakoff-Mayber A (1967) Effect of Mediterranean vegetation on the moisture regime. In WE Sopper, HW Lull, eds. *Forest hydrology*. Pergamon Press. Oxford. pp 291-311

Shalyt MS (1950) Podzemnaja cast' nekotorykh lugovykh, stepnykh i pustynnykh rastenij i fitocenozov. C. I. Travjanistye i polukustarnikovye rastenija i fitocenozy lesnoj (luga) i stepnoj zon. (Belowground parts of some meadow, steppe, and desert plants and plant communities. Part I: Herbaceous plants and

subshrubs and plant communities of forest and steppe zones. In Russian). Trudy Botanicheskogo Instituta im. V.L. Komarova. Akademii nauk SSSR. Seria III. Geobotanika 6: 205-442

Shalyt MS (1952) Podzemnaja cast' nekotorykh lugovykh. stepnykh i pustynnykh rasteniy i fitocenozov. C. 2. Travjanistye. polukustarnikovye i kustarnikovye rastenija i fitocenozy pustynnoj zony. (Belowground parts of some meadow. steppe. and desert plants and plant communities. Part 2: Herbaceous plants. subshrubs. and shrubs. and plant communities of the desert zone. In Russian). Trudy Botanicheskogo Instituta im. V.L. Komarova. Akademii nauk SSSR. Seria III. Geobotanika 8: 71-139

Shankarnarayan KA, Dabadghao PM, Pandey PK, Rai P (1974) Studies on the root system of five principal grasses of Sehima-Dichanthium cover. Annals of Arid Zone 13: 177-186

Sharifi MR, Nilsen ET, Rundel PW (1982) Biomass and net primary production of *Prosopis glandulosa* (Fabaceae) in the Sonoran Desert of California. American Journal of Botany 69: 760-767

Sharma BM (1968) Root systems of some desert plants at Churu. Rajasthan. Indian Forester 94: 240-246

Sharma BM (1984) Ecophysiological studies of *Eleusine indica* (L.) Gaertn. *Sporobulus pyramidalis* P. Beauv. at Ibadan. Nigeria. Journal of Range Management 37: 275-276

Sharma BM, Chivinge AO (1982) Contributions to the ecology of *Dactyloctenium aegyptiacum* (L.) P. Beauv. Journal of Range Management 35: 326-331

Shaver G, Billings DW (1975) Root Production and Root Turnover in a Wet Tundra Ecosystem, Barrow, Alaska. Ecology. 56. 401. 10.2307/1934970.

Sherff E (1912) The vegetation of Skokie Marsh, with special reference to subterranean organs and their interrelationships. Botanical Gazette 53(5):415–435.

Shpak IS (1971) The effects of forests on the water balance of drainage basins. Israel Program for Scientific Translations. Jerusalem

Silva S, Whitford WG, Jarrell WM, Virginia RA (1989) The microarthropod fauna associated with a deep rooted legume. *Prosopis glandulosa*. in the Chihuahuan Desert. Oecologia 7: 330-335

Sin'kovsky LP (1985) Herbs of local flora as used in cultivation. In NT Nechaeva. ed. Improvement of desert ranges in Soviet Central Asia. Vol 4. Harwood Academic Publishers. Chur. Switzerland. pp 203-220

Singh SP (1964) Cover. biomass. and root-shoot habit of *Larrea divaricata* on a selected site in southern New Mexico. Thesis. New Mexico State University. Las Cruces

Singh V (1994) Morphology and pattern of root distribution in *Prosopis cineraria*. *Dalbergia sissoo* and *Albizia lebbek* in an arid region of north-western India. Tropical Ecology 35: 133-146

Skerman PJ, Riveros F (1990) Tropical grasses. Food and Agriculture Organization of the United Nations. Rome

Soethe N, Lehmann J, Engels C (2006) Root morphology and anchorage of six native tree species from a tropical montane forest and an elfin forest in Ecuador. Plant and Soil, 279: 173–185.

Soriano A (1983) Deserts and semi-deserts of Patagonia. In NE West. ed. Temperate deserts and semi-deserts. Vol 5. Elsevier. Amsterdam. pp 423-460

Soriano A, Sala OE (1983) Ecological strategies in a Patagonian arid steppe. Vegetatio 56: 9-15

- Soumaré A, Groot JJR, Koné D. Radersma S (1994) Structure spatiale du système racinaire de deux arbres du Sahel: *Acacia seyal* et *Sclerocarya birrea*. Vol 5. Production Soudano-Sahélienne (PSS). Wageningen
- Specht RL (1957) Dark Island heath (Ninety-mile Plain. South Australia) III. The root system. Australian Journal of Botany 5: 103-114
- Specht RL (1957) Dark Island heath (Ninety-mile Plain. South Australia) IV. Soil moisture patterns produced by rainfall interception and stemflow. Australian Journal of Botany 5: 137-150
- Spence LE (1937) Root studies of important range plants of the Boise River watershed. Journal of Forestry 35: 747-754
- Sperry TM (1935) Root systems in Illinois prairie. Ecology 16: 178-202
- Sprackling JA, Read RA (1979) Tree root systems in eastern Nebraska. Nebraska Conservation Bulletin 37: 73
- Steele SJ, Gower ST, Vogel JG, Norman JM (1997) Root mass. net primary production and turnover in aspen. jack pine and black spruce forests in Saskatchewan and Manitoba. Canada. Tree Physiology 17: 577-587
- Stocker O (1928) Der Wasserhaushalt ägyptischer Wüsten- und Salzpflanzen vom Standpunkt einer experimentellen und vergleichenden Pflanzengeographie aus. Botanische Abhandlungen 13: 200
- Stone E.L, Kalisz PJ (1991) On the maximum extent of tree roots. Forest Ecology and Management. 46(1–2). 59–102. [http://doi.org/10.1016/0378-1127\(91\)90245-Q](http://doi.org/10.1016/0378-1127(91)90245-Q)
- Strong WL, La Roi GH (1983) Rooting depths and successional development of selected boreal forest communities. Canadian Journal of Forest Research 13: 577-588
- Sturges DL (1977) Soil water withdrawal and root characteristics of big sagebrush. American Midland Naturalist 98: 257-273
- Sveshnikova VM (1979) Dominanty Kasakhstanskikh stepej. (Dominant plants of the Kazakhstan steppe In Russian). Nauka. Leningrad
- Tabler RD (1964) The root system of *Artemisia tridentata* at 9.500 feet in Wyoming. Ecology 45: 633-636
- Tadmor NH, Orshan G, Rawitz E (1962) Habitat analysis in the Negev Desert of Israel. Bulletin of the Research Council of Israel 11D: 148-173
- Thomas CM, Davis SD (1989) Recovery patterns of three chaparral shrub species after wildfire. Oecologia 80:309-320.
- Thomas FM (2000) Vertical rooting patterns of mature *Quercus* trees growing on different soil types in northern Germany. Plant Ecology 147: 95-103
- Thomas HH (1921) Some observations on plants in the Libyan desert. Journal of Ecology 9: 75-89
- Thomas WD (1980) Characteristics of root systems: California oaks. In TR Plumb. ed. Proceedings of the symposium on the ecology. management. and utilization of California oaks; 1979 June 26-28; Claremont. CA. Vol PSW-44. U.S. Department of Agriculture. Forest Service. Pacific Southwest Forest and Range Experiment Station. Berkeley. CA. pp 178-179

Tierney GD, Foxx TS (1982) Floristic composition and plant succession on near-surface radioactive waste disposal facilities in the Los Alamos National Laboratory. Vol LA-9212-MS. Los Alamos National Laboratory. Los Alamos. New Mexico

Tierney GD, Foxx TS (1987) Rooting lengths of plants of Los Alamos National Laboratory lands. Vol LA-10865-MS. Los Alamos National Laboratory. Los Alamos. New Mexico

Toky OP, Bisht RP (1992) Observations on the rooting patterns of some agroforestry trees in an arid region of north-western India. Agroforestry Systems 18: 245-263

Tolstead WL (1942) Vegetation of the northern part of Cherry County. Nebraska. Ecological Monographs 12: 255-292

Tryon PR, Chapin FS, III (1983) Temperature control over root growth and root biomass in taiga forest trees. Canadian Journal of Forest Research 13: 827-833

Turner GT, Costello DF (1942) Ecological aspects of the pricklypear problem in eastern Colorado and Wyoming. Ecology 23: 419-426

Upendra S, Good ER (1993) Vertical root distribution in relation to soil properties in New Jersey Pinelands forests. Plant and Soil. 150. 87-97. 10.1007/BF00779179.

USDA, NRCS. 2021. The PLANTS Database (<http://plants.usda.gov>). Greensboro, NC, USA: National Plant Data Team [accessed 10 September 2021].

Van Donselaar-Ten Bokkel Huinink WAE (1966) Structure. root systems and periodicity of savanna plants and vegetations in northern Surinam. Wentia 17: 1-162

Vandenbeldt RJ (1991) Rooting systems of western and southern African *Faidherbia albida* (Del.) A. Chev. (syn. *Acacia albida* Del.) - a comparative analysis with biogeographic implications. Agroforestry Systems 14: 233-244

Villares JB, Tundisi A, Becker M (1953) The subterranean system of colonial grass (Guinea grass) in various soils of the State of São Paulo. Brazil. Journal of Range Management 6: 248-254

Wallace A, Romney EM (1972) Radioecology and ecophysiology of desert plants at the Nevada test site. Vol TID-25954. U.S. Atomic Energy Commission. Washington. D.C.

Walter H (1939) Grasland. Savanne und Busch der arideren Teile Afrikas in ihrer ökologischen Bedingtheit. Jahrbücher für wissenschaftliche Botanik 87: 750-860

Walter H (1963) The water supply of desert plants. In AJ Rutter. FH Whitehead. eds. The water relations of plants. Vol No. 3. John Wiley & Sons. New York. pp 199-205

Walter H, Breckle S-W (1984) Spezielle Ökologie der tropischen und subtropischen Zonen. Vol Band 2. Gustav-Fischer Verlag. Stuttgart. Jena

Walter H, Breckle S-W (1986) Ecological systems of the geobiosphere 2: Tropical and subtropical zonobiomes. Springer-Verlag. Berlin

Wang P, Mommer L, Ruijven JV, Berendse F, Maximov TC, Heijmans MM (2016). Seasonal changes and vertical distribution of root standing biomass of graminoids and shrubs at a Siberian tundra site. Plant and Soil, 407, 55-65.

- Weaver JE (1917) A study of the vegetation of southeastern Washington and adjacent Idaho. University Studies of the University of Nebraska 17: 1-133
- Weaver JE (1919) The ecological relations of roots. Vol No. 286. Carnegie Institution of Washington. Washington. D.C.
- Weaver JE (1920) Root development in the grassland formation. Vol No. 292. Carnegie Institution of Washington. Washington. D. C.
- Weaver JE (1954) North American prairie. Johnsen Publishing Company. Lincoln. Nebraska
- Weaver JE (1958) Summary and interpretation of underground development in natural grassland communities. Ecological Monographs 28: 55-78
- Weaver JE, Albertson FW (1956) Grasslands of the Great Plains. Johnsen Publishing Company. Lincoln. Nebraska
- Weaver JE, Darland RW (1949) Soil-root relationships of certain native grasses in various soil types. Ecological Monographs 19: 303-338
- Weaver JE, Kramer J (1932) Root system of *Quercus macrocarpa* in relation to the invasion of prairie. Botanical Gazette 94: 51-85
- Westman WE, Rogers RW (1977) Biomass and structure of a subtropical eucalypt forest. North Stradbroke Island. Australian Journal of Botany 25: 171-191
- Williams K, Hoobs RJ (1989) Control of shrub establishment by springtime soil water availability in an annual grassland. Oecologia 81: 62-66
- Wilson RC (1972) *Abronia*: I. Distribution, ecology and habit of nine species of *Abronia* found in California. Aliso 7: 421-437
- Wright CD (1928) An ecological study of *Baccharis pilularis*. M.S. thesis. University of California. Berkeley
- Wright SJ, Machado JL, Mulkey SS, Smith AP (1992) Drought acclimation among tropical forest shrubs (*Psychotria*, Rubiaceae). Oecologia 89: 457-463
- Wyatt JW, Dollhopf DJ, Schafer WM (1980) Root distribution in 1- to 48 year-old strip-mine spoils in southeastern Montana. Journal of Range Management 33: 101-104
- Yeager AF (1935) Root systems of certain trees and shrubs grown on prairie soil. Journal of Agricultural Research 51: 1085-1092
- Yeaton RI, Travis J, Gilinsky E (1977) Competition and spacing in plant communities: the Arizona upland association. Journal of Ecology 65: 587-595
- Yoshisuke N, Brown J (1972) Mathematical Modeling and Validation of the Thermal Regimes in Tundra Soils, Barrow, Alaska, Arctic and Alpine Research, 4:1, 19-38
- Zohary M (1961) On hydro-ecological relations of the Near East desert vegetation. UNESCO Arid Zone Research 16: 199-212
- Zverev NE, Seiidova RD (1990) Underground mass of shrub and subshrub plants of the Karakum. Problems of Desert Development 1990(1): 49-54