

New Scientist invites you to experience 'extreme nature' on

4TH - 13TH DECEMBER 2006
10 DAY CRUISE - \$3395*

THE ULTIMATE Galapagos IN-DEPTH VOYAGE

NewScientist.com

26 September 2006

HOME | NEWS | EXPLORE BY SUBJECT | LAST WORD | SUBSCRIBE | SEARCH | ARCHIVE | RSS | JOBS

SEARCH

Email Sign-up
Subscribe to Magazine
Customer Service

New SUMMER OFFER FOR 4 EXTRA FREE ISSUES

NewScientist.com podcast

NS Full Access

BREAKING NEWS**LATEST HEADLINES**

[Pool knowledge to find the origins of language](#)
[Virtual face-ageing may help find missing persons](#)
[Predicting the impact of wild habitat loss](#)
[Scanner recognises hidden knives and guns](#)
[Global warming nears a million-year high](#)
[Ice the size of Texas melts in one year](#)
[Ageing may be a lifestyle 'choice' for bacteria](#)
[First zero-gravity surgery set to be performed](#)
[ALL LATEST NEWS](#)

PRINT EDITION[Subscribe](#)

- Current issue
- Archive
- NS Full Access

JOB**JOB OF THE WEEK****Postdoctoral Fellow****SUBSCRIPTIONS****4 FREE ISSUES**[Click here](#)

[Subscribe](#)
[Renew](#)
[Change address](#)

Give a NewScientist Gift Subscription *
Save 70% *
[Click here *](#)

The World's No.1 Science & Technology News Service**Predicting the impact of wild habitat loss**

15:59 26 September 2006
 NewScientist.com news service
 John Pickrell

Habitat loss is one of the greatest threats many species face, but scientists have struggled to predict the effects of, for example, rainforest loss, due to the sheer numbers of species involved.

Researchers have now focused on a single self-contained ecosystem – comprised simply of a carnivorous plant and the species that inhabit it – and used this to create a model that could apply to much larger systems.

The study is the first wild experiment to test how the interactions of predators and prey can drive decline when habitats contract, the researchers claim. "To actively manipulate an entire food web is pretty difficult outside the laboratory," explains Nicholas Gotelli, at the University of Vermont in Burlington, US.

To overcome that problem, Gotelli and colleague Aaron Ellison turned to the northern pitcher plant (*Sarracenia purpurea*), a carnivorous species found in the bogs of southern Canada and the eastern US. The plant produces up to 12 fat, tubular shaped leaves each year that fill with water and support an aquatic community.

Food webs

The pitcher plants use nectar, and bright colouration to attract ants and flies into the leaves, which fall into the water and drown. They form the basis of a complex food web, involving water-bound midge and fly larvae that eat them. Other tiny species, such as bacteria, protozoa, and rotifers, eat the leftovers – and also serve as food for the larvae.

The complexity of this food web is comparable to that of food webs of much larger species, says Gotelli.

Working with 50 pitcher plants in a Vermont bog, the researchers reduced or added water, to stimulate habitat loss, and in some cases removed the midge and fly larvae which act as "top predators" in these miniature ecosystems. Top predators are often the first to vanish when habitats contract, as they need the largest areas to survive.

Complex interactions

Gotelli and Ellison compared a range of computer models of how species are predicted to decline after habitat loss with the pair's actual results of how species had declined. Models that took into account the complex interactions of species were much better at predicting actual declines, than those that made predictions based on habitat size alone, they confirmed.

"If we're going to understand the effects of habitat loss on natural communities, we're going to have to do more than study a single species at a time," says Gotelli.

He plans to use his model to study the effects of climate change on the ecosystem, first varying the amounts of precipitation.

Looking at the total interactions of predators and prey is important in formulating methods to save species from decline, he adds.

Journal reference: *PLoS Biology* (DOI: 10.1371/journal.pbio.0040324)

[PRINT](#) [SEND](#) [RSS FEEDS](#) [SYNDICATE](#)

Tools

[digg this](#)
[reddit](#) [SUBMIT](#)
[ADD MY YAHOO!](#)
[NEWS VINE](#)
[DEL.icio.us](#)

[Enlarge image](#)

The carnivorous northern pitcher plant hosts an entire miniature community of species (Image: Aaron M Ellison)

Related Articles

[When a deathtrap becomes a free lunch](#)
 24 December 2005

[All life is here - Small, stagnant pools teeming with wriggling larvae are helping to](#)
 15 March 1997

[EU's butterfly havens are vanishing](#)
 17 March 2006

[Search New Scientist](#)
[Contact us](#)

Web Links

[PloS Biology](#)
[Nicholas Gotelli](#)
[Endangered Species, New Scientist special report](#)

SUBSCRIBER LOGIN

username:
 password:
 Your login is case-sensitive

[Log in](#)

- Forgotten your password?
- [Subscriber? Register now!](#)
- [Subscribe now](#)
- [Institutional Subscribers](#)
- [Athens login](#)

NewScientist

New Scientist invites you to experience 'extreme nature' on

THE ULTIMATE Galapagos IN-DEPTH VOYAGE

4TH - 13TH DECEMBER 2006
 10 DAY CRUISE - \$3395*



Peregrine

GET 51 ISSUES FULL ONLINE ACCESS AND OVER 70% OFF



SPONSORED LINKS

[Car Insurance by A Quote](#)
[Van Insurance by A Quote](#)
[Bike Insurance by A Quote](#)

[Home Mortgage Loans](#)
[Movers in Your area](#)
[Find Moving Companies](#)
[Real Estate](#)
[International Movers](#)

[Contact us about links](#)



- For exclusive news and expert analysis every week [subscribe](#) to New Scientist Print Edition
- For what's in New Scientist magazine this week see [contents](#)
- [Search](#) all stories
- [Contact us](#) about this story
- [Sign up](#) for our free newsletter

[Subscribe](#) [Contact Us](#) [FAQ / Help](#) [Advertise](#) [Disclaimer](#) [Terms and Conditions](#) [Cookies](#) [Privacy Policy](#) [Open Source](#) [Site Map](#)
[About NewScientist.com](#) [About New Scientist magazine](#) [Publisher Blog](#) © Copyright Reed Business Information Ltd. [Vacancies](#)

[Home](#)

