



innovations report

Forum für Wissenschaft, Industrie und Wirtschaft

Hauptsponsoren: **SIEMENS** **n-tv** **Postbank**

Datenbankrecherche: Fachgebiet (optional):

[Home](#) [Über uns](#) [English](#)

FACHGEBIETE **SONDERTHEMEN** **FORSCHUNG** **B2B BEREICH** **JOB & KARRIERE** **SERVICE**

NACHRICHTEN & BERICHTE

- Agrar- Forstwissenschaften
- Architektur Bauwesen
- Automotive
- Biowissenschaften Chemie**
- Energie und Elektrotechnik
- Geowissenschaften
- Gesellschaftswissenschaften
- Informationstechnologie
- Interdisziplinäre Forschung
- Kommunikation Medien
- Maschinenbau
- Materialwissenschaften
- Medizintechnik
- Medizin Gesundheit
- Ökologie Umwelt- Naturschutz
- Physik Astronomie
- Studien Analysen
- Verfahrenstechnologie
- Verkehr Logistik
- Wirtschaft Finanzen

Weitere Förderer



Ads by Google

Ant Habitat

Bug Habitat

Home → Fachgebiete → Biowissenschaften Chemie → Nachricht

Predicting species abundance in the face of habitat loss

26.09.2006

[> nächste Meldung >](#)

Habitat loss poses the greatest threat to the survival of a species, and often precipitates the demise of top predators and wide-ranging animals, like the Siberian tiger and the orangutan.



Any hope of recovering such critically endangered species depends on understanding what drives changes in population size following habitat contraction. In a new study published in PLoS Biology, Nicholas Gotelli and Aaron Ellison test the relative contributions of habitat contraction, keystone species effects, and food-web interactions on species abundance, and provide experimental evidence that trophic interactions exert a dominant effect. Until now, direct evidence that trophic interactions play such an important role has been lacking, in part because manipulating an intact food web has proven experimentally intractable, and in part because these different modeling frameworks have not been explicitly compared.

Ads by Goooooogle

Gotelli and Ellison overcame such technical limitations by using the carnivorous pitcher plant (*Sarracenia purpurea*) and its associated food web as a model for studying what regulates abundance in shrinking habitats. Every year, the pitcher plant, found in bogs and swamps throughout southern Canada and the eastern United States, grows six to 12 tubular leaves that collect enough water to support an entire aquatic food web. The pitcher plant food web starts with ants, flies, and other arthropods unlucky enough to fall into its trap. Midges and sarcophagid fly larvae "shred" and chew on the hapless insect. This shredded detritus is further broken down by bacteria, which in turn are consumed by protozoa, rotifers, and mites. Pitcher plant mosquito larvae feed on bacteria, protozoa, and rotifers. Older, larger sarcophagid fly larvae also feed on rotifers as well as on younger, smaller mosquito larvae.

Working with 50 pitcher plants in a bog in Vermont, Gotelli and Ellison subjected the plants to one of five experimental treatments, in which they manipulated habitat size (by changing the volume of water in the leaves), simplified the trophic structure (by removing the top trophic level—larvae of the dipterans fly, midge, and mosquito), did some combination of the two, or none of the above (the control condition). Dipteran larvae and water were measured as each treatment was maintained; both were replaced in the control condition and more water was added in the habitat expansion treatment. These treatments mimic the kinds of changes that occur in nature as habitat area shrinks and top predators disappear from communities.

The best predictors of abundance were models that incorporated trophic structure—including the "mosquito keystone model." This model accurately reflected the pitcher plant food web, with mosquito larvae preying on rotifers, and sarcophagid flies preying on mosquito larvae. "Bottom-up" food web models (in



Aktuell

- Airline-Boss Branson will Schadstoff-Ausstoß von Jets reduzieren**
28.09.2006 | Ökologie Umwelt-Naturschutz
- Bundesregierung baut technologische Kooperation mit Vietnam aus**
28.09.2006 | Bildung Wissenschaft
- CBEX und Deutsche Börse unterzeichnen Kooperationsvereinbarung**



which links flow from prey to predator) predicted that changes in bacteria population size influence protozoa abundances, which in turn affect mosquito numbers, and that changes in bacteria abundance also affect mite numbers, which impact rotifer abundance. This scenario lends support to the model of a Sarracenia food web in which each link in the chain performs a specialized service in breaking down the arthropod prey that is used by the next species in the processing chain.

With over 200 million acres of the world's forestlands destroyed in the 1990s alone, and an estimated 40% increase in the human population by 2050, a growing number of species will be forced to cope with shrinking habitat. Instead of trying to determine how individual species might respond to habitat loss, Gotelli and Ellison argue that incorporating trophic structure into ecological models may yield more-accurate predictions of species abundance—a critical component of species restoration strategies.

Citation: Gotelli NJ, Ellison AM (2006) Food-web models predict species abundances in response to habitat change. PLoS Biol 4(10): e324. DOI: 10.1371/journal.pbio.0040324.

Andrew Hyde | Quelle: alphagalileo
 Weitere Informationen: dx.doi.org/10.1371/journal.pbio.0040324
 www.plosbiology.org

> nächste Meldung >

[Ads by Goooooogle](#) [Advertise on this site](#)

[Mosquito Ringtone](#)
 Enter Your Number To Get The Mosquito Tone Plus 10 Bonus Tones!
 www.Ringtones.com/Mosquito

[Got Asian Tiger Mosquito?](#)
 Mosquito Magnet Lurex specially designed for Asian Tiger Mosquito
 www.mosquitomagnet.com



☑ Top ✉ Artikel versenden 🖨 drucken

