

Reburn severity in managed and unmanaged vegetation in a large wildfire

Jonathan R. Thompson, Thomas A. Spies, and Lisa M. Ganio

PNAS published online Jun 11, 2007;

doi:10.1073/pnas.0700229104

This information is current as of June 2007.

Supplementary Material

Supplementary material can be found at:
www.pnas.org/cgi/content/full/0700229104/DC1

This article has been cited by other articles:
www.pnas.org/otherarticles

E-mail Alerts

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

Rights & Permissions

To reproduce this article in part (figures, tables) or in entirety, see:
www.pnas.org/misc/rightperm.shtml

Reprints

To order reprints, see:
www.pnas.org/misc/reprints.shtml

Notes:

Reburn severity in managed and unmanaged vegetation in a large wildfire

Jonathan R. Thompson*[†], Thomas A. Spies[‡], and Lisa M. Ganio*

*Department of Forest Science, Oregon State University, Corvallis, OR 97331; and [†]Pacific Northwest Research Station, U.S. Department of Agriculture Forest Service, Corvallis, OR 97331

Edited by Ruth S. DeFries, University of Maryland, College Park, MD, and approved April 26, 2007 (received for review January 10, 2007)

Debate over the influence of postwildfire management on future fire severity is occurring in the absence of empirical studies. We used satellite data, government agency records, and aerial photography to examine a forest landscape in southwest Oregon that burned in 1987 and then was subject, in part, to salvage-logging and conifer planting before it reburned during the 2002 Biscuit Fire. Areas that burned severely in 1987 tended to reburn at high severity in 2002, after controlling for the influence of several topographical and biophysical covariates. Areas unaffected by the initial fire tended to burn at the lowest severities in 2002. Areas that were salvage-logged and planted after the initial fire burned more severely than comparable unmanaged areas, suggesting that fuel conditions in conifer plantations can increase fire severity despite removal of large woody fuels.

public land management | salvage-logging | Biscuit Fire | Landsat | landscape ecology

Large wildfires are increasingly common in western North America (1). Changing climate patterns and the legacy of fire suppression within fire-prone forests suggest that this trend will continue. Postfire management is, therefore, a growing concern for public land managers. Although it has been customary to salvage-log fire-killed trees and plant seedlings after large wildfires, there is a mounting debate regarding the practice (2–4). There are several reasons one might choose this management system, including recouping economic losses through timber sales and ensuring the reestablishment of desirable tree species. Another common justification for this approach has been a perceived reduction in future fire risk associated with the removal of dead wood (2, 5–7). The threat of severe reburns is real but not well understood (4). For example, Oregon's Tillamook burns of the 1930s, 1940s, and 1950s consisted of one large fire followed by three reburns 6, 12, and 18 years later. In sum, these fires burned more than 135,000 hectares. The threat of reburns motivates public land managers to construct fuel-breaks and to salvage-log to hedge against the risks of future fire (6). Recent studies have found, however, that salvage-logging can increase surface fuels available to fires above prelogging levels by transferring unmerchantable material to the forest floor, suggesting that this postfire management practice might actually increase fire risk for a time (3, 8). Until now, no study has quantified how recent fire history and postfire management actually affects the severity of a large wildfire (4).

The 2002 Biscuit Fire was among the largest forest fires in modern United States history, encompassing >200,000 hectares primarily within the Rogue-Siskiyou National Forest (RSNF) in southwest Oregon. In the years following, the Biscuit Fire has been a catalyst for national debate regarding forest management in the aftermath of wildfires on public land. This debate is taking place in the absence of empirical research on how future wildfire severity is associated with past wildfires and how postfire forest management alters future fire severity (4). We analyzed burn severity patterns within 18,000 hectares of the Biscuit Fire that burned 15 years earlier during the 1987 Silver Fire. Both fires burned heterogeneously, creating mosaics of live and dead trees

in variably sized patches. In the 3 years following the Silver Fire, >800 hectares were salvage-logged and planted with conifers. The arrangement of these disturbances presented a unique opportunity to address two important research questions. First, was severity in the Biscuit Fire associated with severity in the Silver Fire in unmanaged areas? Second, did areas that were salvage-logged and planted with conifers after the Silver Fire burn more or less severely in the Biscuit Fire than comparable unmanaged areas?

With regard to the first question, hereafter referred to as “the reburn question,” a negative correlation between Biscuit and Silver Fire severity is plausible if the forests that burned severely in 1987 had less remaining fuel to support the Biscuit Fire in 2002, or if regenerating young forests did not effectively carry fire. This relationship has been observed in lodgepole pine ecosystems (9–11). An alternate hypothesis is that Biscuit Fire severity would be positively correlated with Silver Fire severity. This would occur if areas of higher Silver Fire severity had greater accumulations of fire-killed trees and vegetative growth available as fuel to the Biscuit Fire. This scenario is assumed to have influenced forest dynamics in more mesic forests of the Pacific Northwest (12). Finally, there may be no discernible association between the severity patterns of the two fires. Many independent factors influence fire severity, including weather, topography, fuel, landscape structure, and fire suppression. Any of these could overwhelm the signal from the legacy of the Silver Fire.

The second question, hereafter referred to as “the salvage-plant question,” also has several plausible outcomes. The hypothesis that salvage-logging followed by planting conifers can reduce future fire severity is widely held and rests on the assumption that removing dead trees reduces fuel loads, and planting conifers and controlling competing vegetation hastens the return of fire-resistant forests (2, 5–7). An alternative hypothesis is that salvage-logging plus plantation creation exacerbates future fire severity. No studies have measured fire severity following salvage-logging, but it is known that it can increase available fine and coarse fuel loads if no fuel treatments are conducted (3, 8). In addition, several studies have documented high-severity fire within young conifer plantations, where surface fuels can be fine, homogeneous, and continuous (13–15).

Our study area is within the Siskiyou Mountains in southwest Oregon's mixed-conifer and mixed-evergreen hardwood zones

Author contributions: J.R.T., T.A.S., and L.M.G. designed research; J.R.T. performed research; J.R.T. and L.M.G. analyzed data; and J.R.T. and T.A.S. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Abbreviations: RSNF, Rogue-Siskiyou National Forest; dNBR, differenced normalized burn ratio; TM, Thematic Mapper; PAG, plant association group.

[†]To whom correspondence should be addressed. E-mail: jonathan.thompson@oregonstate.edu.

This article contains supporting information online at www.pnas.org/cgi/content/full/0700229104/DC1.

© 2007 by The National Academy of Sciences of the USA




Fig. 3. Ninety-five percent confidence intervals for Biscuit Fire dNBR, a Landsat-derived burn severity metric. Estimates and confidence intervals compare regions burned at high severity in the 1987 Silver Fire that were unmanaged to areas that were salvage-logged and planted after the Silver Fire. Means were calculated at the multivariate centroid of the covariates: slope, elevation, topographic position, Silver Fire severity, day-of-burn, and 1986 greenness (see Table 2). Means were similar across the four PAGs. In this landscape, the Tanoak group is found on wetter sites in the western portion of the study area, whereas the Tanoak-Canyon live oak group is found on dryer, inland sites; the Douglas-fir group is found on relatively dry sites; and the White-fir group is found at somewhat higher elevation wetter sites (20). Colors correspond to Biscuit Fire severity calibrated through comparison with aerial photography: blue, <10% canopy scorch; green, 10–50% canopy scorch; red, >50% canopy scorch.

Discussion and Conclusion

No previous study has compared fire severity in plantations and naturally regenerated vegetation of similar ages. Our findings are consistent with studies that show that site history influences fire severity (15, 24, 25), and with studies that have found an association of high-severity fire with conifer plantations (13–15). Our limited knowledge of the fuel characteristics at the time of the Biscuit Fire prevents us from separating the effects of logging and planting. The relative influence of these management actions on burn severity would vary over time: the influence of dead fuels and harvest debris would diminish as they decayed (3) and the influence of live vegetation would increase as it developed. The patterns we observed apply to the particular conditions and history of post-Silver Fire management; they could change with shorter or longer intervals between fires.

The Biscuit Fire tended to burn at relatively high severity in young naturally regenerated stands and even more severely in young conifer plantations of comparable age and fire history. This suggests that young forests, whether naturally or artificially regenerated, may be vulnerable to positive feedback cycles of high severity fire, creating more early-successional vegetation and delaying or precluding the return of historical mature-forest composition and structure. Although patches of high-severity fire and reburns are a normal part of the mixed-severity fire regime within this forest type (12, 26), increasing occurrence of wildfire driven by climate warming in this region (1) may lead to increases in the prevalence of sclerophyllous species, which are adapted to frequent severe fires (12, 16, 20).

Our findings are inconsistent with the hypothesis that this particular postfire management system reduces the risk of high-severity fire in a reburn occurring 15 years after the original

fire. The logging component of this system is often considered a fuel-reduction treatment (2, 5–7). However, the large-diameter fuels removed during harvest do not readily carry wildland fire (12, 27). Thus, logging may not reduce available fuels. In fact, harvesting fire-killed trees may increase available surface fuels by transferring unmerchantable material, such as tops, branches, and broken boles to the ground immediately after harvest (3, 8). This effect may be mitigated as logging slash decays, or through fuel reduction methods, such as broadcast-burning (15). Records of site preparation and their effectiveness in reducing fuels in the plantations are incomplete; however, at least 17 of the 44 plantations are reported as “broadcast-burned.” In a separate analysis, we found that these 17 plantations also burned with higher severities than comparable unmanaged stands. The planting component of the system is intended to promote long-term regrowth of conifer trees, but it also creates dense or continuous fuels that are at elevated risk of high severity fire (14). It should be noted, however, that many of the plantations examined in this analysis had lower conifer densities and a larger component of shrubs and hardwoods than would be found in typical intensively managed plantations of the same age (11–14 years). Our analysis could not measure any details regarding differences in preburn structure and composition between the natural and artificially regenerated stands. Nonetheless, the naturally regenerated areas received no site preparation or planting; therefore, they likely contained a more diverse arrangement of young vegetation and open gaps (20, 28). Although these naturally regenerated areas also supported relatively high-severity fire, abrupt changes in fuel profiles, which can slow fire spread (12), may have reduced the average burn severity.

We currently lack general conceptual models or simulation models that can help us understand the effects of salvage-logging on fire severity over large landscapes and long time frames. As our work indicates, research needs to consider all of the components of postfire management systems, individually and together. Thus far, the few studies that have examined reburn potential in salvage-logged sites have emphasized the dead woody fuel transferred to the surface during harvest. But logging slash is only part of the fire risk story, and it may not be the most important after a few years. On public land, salvage-logging is almost always followed by conifer planting, even when the objective is ecological recovery, such as expediting the return of old-growth forests. We are currently unable to examine the short- and long-term tradeoffs associated with different postfire management systems. For example, we do not know how the apparent difference in fire hazard between plantations and natural stands that we observed at 15 years varies over time and whether this short-term risk is balanced in any way by longer term benefits in terms of stand development and reduced fire risk. However, the available evidence suggests that the combined influence from a pulse input of surface fuels resulting from salvage-logging (3, 8) followed by the establishment of uniform young plantations may increase susceptibility to severe reburns in the early stages of forest development.

Managers may have few options to reduce the risk of high-severity fire within areas that have recently burned severely. Typical fuel treatments, such as thinning, do not have much effect on fire risk in young forests (14). Reducing connectivity of surface fuels at landscape scales is likely the only way to decrease the size and severity of reburns until vertical diversification and fire resistance is achieved (29). The decision to salvage-log and plant, or not, after fire depends on a number of management considerations including risk of future high-severity fire, reducing hazards to fire fighters, timber revenue, and conservation of biodiversity. Further research, especially controlled experiments, is clearly needed to help managers understand tradeoffs. Given the difficulty of conducting experiments with large wild-

from 600 to 1,100 trees per hectare. In addition to the logged and planted areas used in the analysis, we distinguished ≈ 250 hectares that were harvested in part or in full after the Silver Fire but were either not planted or the planted conifers were not certified “free to grow.” We excluded these later areas from all our analyses because their history was too uncertain and variable to accurately characterize. All management polygons were provided by the RSNF and were edited in a geographic information system (GIS) by using a 1-m Digital Orthoquad acquired in 1994 to better fit the perimeters of harvest units. Areas logged and planted before the Silver Fire were excluded from all analyses. The bulk of the remaining 17,000 hectares in the northern Silver Fire perimeter were not harvested. However, some small unknown proportion was selectively logged but not planted; no records describing management actions are known to exist and we could not see evidence of them in careful analysis of the Digital Orthoquads. It is likely that only the largest most valuable trees were extracted from these sites via helicopter. The unrecorded, lightly salvaged sites make up no more than 10% of the study area and are included in the population of unmanaged sites.

Sampling, Variable Selection, Model Fitting, and Hypothesis Testing.

All sampling and data extraction was done in a geographic information system in which all of the variables were converted to 29-m raster maps. The sample universe for the reburn question was the region within the northern Silver Fire but outside any management areas. All known management units dating back >50 years were excluded from reburn analysis. Sample locations were determined by randomly selecting locations for 381 points with the constraint that they be separated by at least 300 m to reduce spatial dependence. Sample values were calculated as the mean value of the closest nine contiguous pixels to the sample location (sample unit area, ≈ 0.75 hectares). When sampling for the salvage-plant question, we constrained the sample universe to the areas within the northern Silver Fire perimeter, areas within PAGs that contained

management units and either burned at high severity in 1987 and received no or minimal postfire management, or areas that burned at high severity and were clearcut during 1988, 1989, or 1990, and then planted in the years following and certified by the RSNF as an established conifer plantation. All other management units were excluded. We sampled 292 random locations (225 unmanaged and 67 managed) separated from each other by at least 300 m. Any plantation that did not include at least one sample during the initial sample selection had a sample randomly located within it. Data were extracted from all of the raster maps as described above.

Statistical analysis was completed in the computing software R (37). Empirical variogram models indicated spatial autocorrelation of Biscuit dNBR data; a spherical theoretical variogram model best described the autocorrelation. This spatial dependence precluded the use of ordinary least squares regression. Instead, for model selection and hypothesis testing, we used generalized least squares regression to fit linear models of predictor variables to Biscuit dNBR data. Generalized least squares models allow residuals to have a nonstandard covariance structure (38); we used a spherical spatial correlation structure. The modeled variogram from the reburn data had a range of 2,314 m, and the nugget:sill was 0.387. The modeled variogram from the salvage-plant data had a range of 1,096 m, and the nugget:sill was 0.399. Akaike information criteria were used for model selection (39, 40). For both questions, we evaluated Akaike information criteria scores from ≈ 100 *a priori* candidate covariate models, including global models containing all non-correlated predictor variables and null models that contained none. All candidate models contained plausible combinations of predictor variables based on what is known about fire behavior and from previous studies of burn severity patterns on other fires [supporting information (SI) Table 1].

We thank K. Olsen, K. Pierce, T. Link, and the RSNF for technical assistance. T. Atzet, T. Link, P. Harcomb, W. Cohen, M. Huso, J. Fontaine, D. Donato, H. Salwasser, K. N. Johnson, T. Sensenig, M. Wimberly, J. Bailey, and two anonymous reviewers provided helpful comments on earlier drafts. This research was funded by the Joint Fire Science Program.

1. Westerling AL, Hidalgo HG, Cayan DR, Swetnam TW (2006) *Science* 313:940–943.
2. Gorte RL (2006) *Forest Fire/Wildfire Protection* (Congressional Research Service, Washington, DC), RL30755, p 30.
3. McIver JD, Ottmar R (2007) *For Ecol Manage* 238:268–279.
4. McIver JD, Starr L (2001) *West J Appl Forestry* 16:159–168.
5. Brown JK, Reinhardt ED, Kramer KA (2003) *Coarse Woody Debris: Managing Benefits and Fire Hazard in the Recovering Forest* (US Department of Agriculture Forest Service Rocky Mountain Research Station, Ogden, UT), RMRS-GTR-105, p 16.
6. Rogue Siskiyou National Forest (2004) *Biscuit Fire Recovery Project, Final Environmental Impact Statement* (US Department of Agriculture Forest Service, Pacific Northwest Region, Medford, OR).
7. Sessions J, Bettinger P, Buckman R, Newton M, Hamann AJ (2004) *J Forestry* 102:38–45.
8. Donato DC, Fontaine JB, Campbell JL, Robinson WD, Kauffman JB, Law BE (2006) *Science* 311:352.
9. Turner MG, Romme WH, Gardner RH (1999) *Int J Wildland Fire* 9:21–36.
10. Despain DG, Sellers RE (1977) *Western Wildlands* 21–24.
11. Romme WH (1982) *Ecol Monogr* 52:199–221.
12. Agee JK (1993) *Fire Ecology of Pacific Northwest Forests* (Island, Washington, DC).
13. Odion DC, Frost EJ, Strittholt JR, Jiang H, Dellasala DA, Moritz MA (2004) *Conserv Biol* 18:927–936.
14. Stephens SL, Moghaddas JJ (2005) *Biol Conserv* 125:369–379.
15. Weatherspoon CP, Skinner CN (1995) *Forest Sci* 41:430–451.
16. Franklin JF, Dyrness CT (1988) *Natural Vegetation of Oregon and Washington* (Oregon State Univ Press, Corvallis).
17. Lutes DC, Keane JF, Caratti CH, Key CH, Benson NC, Gangi LJ (2004) *FIREMON: Fire Effects Monitoring and Inventory System* (US Department of Agriculture Forest Service, Rocky Mountain Research Station, Ogden, UT), Vol RMRS-GTR-164-CD, p 400.
18. Brewer CK, Winne JC, Redmond RL, Opitz DW, Mangrich MV (2005) *Photogrammetric Eng Remote Sensing* 71:1311–1320.
19. Miller JD, Yool SR (2002) *Remote Sensing Environ* 82:481–496.
20. Hobbs SD, Tesch SD, Owston PW, Stewart RE, Tappeiner JC, Wells G (1992) *Reforestation Practices in Southwestern Oregon and Northern California* (Forest Research Laboratory, Oregon State Univ, Corvallis, Oregon).
21. Cohen WB, Spies TA, Fiorella M (1995) *Int J Remote Sens* 16:721–746.
22. Crist EP, Cicone RC (1984) *IEEE Trans Geosci Remote Sens* 22:256–263.
23. Taylor AH, Skinner CN (2003) *Ecol Appl* 13:704–719.
24. Raymond CL, Peterson DL (2005) *Can J For Res* 35:2981–2995.
25. Finney MA, McHugh C, Grenfell IC (2005) *Can J For Res* 35:1714–1722.
26. Taylor AH, Skinner CN (1998) *For Ecol Manage* 111:285–301.
27. Anderson HE (1982) *Aids to Determining Fuel Models for Estimating Fire Behavior* (US Department of Agriculture Forest Service Intermountain Forest and Range Experiment Station, Ogden, UT), GTR-INT-122, p 22.
28. Shafford JPA, Hibbs DE, Puettmann KJ (2007) *J Forestry* 105:139–146.
29. Agee JK, Bahro B, Finney MA, Omi PN, Sapsis DB, Skinner CN, van Wagtenonk JW, Weatherspoon CP (2000) *Forest Ecol Manage* 127:55–66.
30. Whittaker RH (1960) *Ecol Monogr* 30:279–338.
31. Waring RH, Milner KS, Jolly WM, Phillips L, McWethy D (2006) *Forest Ecol Manage* 228:285–291.
32. Siskiyou National Forest (1988) *Silver Fire Recovery Project, Final Environmental Impact Statement* (US Department of Agriculture Forest Service, Pacific Northwest Region, Grants Pass, OR).
33. Kennedy RE, Cohen WB (2003) *Int J Remote Sens* 24:3467–3490.
34. Chavez PS (1996) *Photogrammetric Eng Remote Sens* 62:1025–1036.
35. Canty MJ, Nielsen AA, Schmidt M (2004) *Remote Sens Environ* 91:441–451.
36. Schroeder TA, Cohen WB, Song S, Canty MJ, Zhiqiang Y (2006) *Remote Sens Environ* 103:16–26.
37. R Development Core Team (2006) R: A Language and Environment for Statistical Computing (R Foundation for Statistical Computing, Vienna).
38. Venables WN, Ripley BD (1997) *Modern Applied Statistics with S-Plus* (Springer, New York).
39. Akaike E (1973) in *Second International Symposium on Information Theory*, eds Petrov BN, Csaki FF (IEEE, Budapest, Hungary), pp 267–281.
40. Burnham KP, Anderson DR (2002) *Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach* (Springer, New York).